

Two complete mitochondrial genomes of the subfamily Chloroperlinae (Plecoptera: Chloroperlidae) and their phylogenetic implications

Rao-Rao Mo^{1,2}, Ying Wang², Jin-Jun Cao², Guo-Quan Wang¹, Wei-Hai Li², Dávid Murányi³

1 *Guangxi Key Laboratory of Agric-Environment and Agric-Products Safety and National Demonstration Center for Experimental Plant Science Education, Agricultural College, Guangxi University, Nanning, Guangxi 530004, China*

2 *Henan International Joint Laboratory of Taxonomy and Systematic Evolution of Insecta, Henan Institute of Science and Technology, Xinxiang, Henan 453003, China*

3 *Department of Zoology, Eszterházy Károly Catholic University, Leányka u. 6, Eger H-3300, Hungary*

<http://zoobank.org/A7F2955E-35FC-400D-8F85-0113D4571680>

Corresponding authors: Guoquan Wang (wangguoquan0@163.com), Weihai Li (lwh7969@163.com)

Received 18 November 2021

Accepted 4 March 2022

Published 20 May 2022

Academic Editors Christiane Weirauch, Klaus-Dieter Klass

Citation: Mo R-R, Wang Y, Cao J-J, Wang G-Q, Li W-H, Murányi D (2022) Two complete mitochondrial genomes of the subfamily Chloroperlinae (Plecoptera: Chloroperlidae) and their phylogenetic implications. Arthropod Systematics & Phylogeny 80: 155–168. <https://doi.org/10.3897/asp.80.e78173>

Abstract

Two new complete mitochondrial genomes (mitogenomes) of the subfamily Chloroperlinae, *Haploperla japonica* Kohno, 1946 and *Sweltsa* sp., were sequenced. The two species showed similar gene order, nucleotide composition, and codon usage. The *Sweltsa* sp. and *H. japonica* mitogenomes were typical circular DNA molecules, with lengths of 15,893 bp and 16,012 bp, respectively. Standard ATN start and TAN stop codons were present in most PCGs. All tRNA genes exhibited the cloverleaf secondary structure typical for metazoans except the *tRNA^{Ser(AGN)}*, which lacked the dihydrouridine arm. In both species, the secondary structure of *lrRNA* contained five structural domains, while the *srRNA* included three domains. The A+T-rich regions contained different repeat regions in each species. Phylogenetic analyses using Bayesian inference (BI) and maximum likelihood methods (ML) showed identical results. The family Chloroperlidae was sister to the Perlodidae. Our analyses inferred relationships between six of the seven Systellognatha families: (((Chloroperlidae + Perlodidae) + Perlidae) + (Styloperlidae + Pteronarcyidae)) + Peltoperlidae.

Keywords

Chloroperlinae, *Haploperla*, mitochondrial genomes, phylogenetics, Plecoptera, *Sweltsa*

1. Introduction

The mitochondrial genome (mitogenome) of metazoans is 14 to 20 kb in size, and usually a circular, double-stranded molecule. It contains 13 protein coding (PCGs), 2 ribosomal RNA (rRNA) and 22 transfer RNA (tRNA) (altogether 37) genes, and an A+T-rich region which is known as the control (CR) or non-coding region (Wolstenholme

1992; Boore 1999; Gissi et al. 2008). The Plecoptera (stoneflies) constitutes an ancient order of hemimetabolous winged insects. Stoneflies include 308 existing genera in 17 families, including approximately 4,000 valid species worldwide (DeWalt et al. 2021). Chloroperlidae is one of the smaller families, with two subfamilies:

Chloroperlinae Okamoto, 1912 and Paraperlinae Ricker, 1943. In the Chloroperlinae, there are three tribes and 206 described species, while in the Paraperlinae, only two genera and six species are recorded (Alexander and Stewart 1999; Judson and Nelson 2012; Chen and Du 2015a; Li et al. 2015a, 2015b). In the present study, *Haploperla japonica* and *Sweltsa* sp. were analysed; both belong to Chloroperlinae. Zwick (2000) proposed a higher classification for the world fauna of Plecoptera, which is the current generally accepted system. However, the positions of four families (Chloroperlidae, Perlidae, Perlodidae, Notonemouridae) were still not clear (Zwick 2000). So far, the phylogenetic position of Plecoptera within the Neoptera, and phylogenetic relationships among some Plecoptera families have remained controversial, and more molecular data are needed to resolve them (Stewart and Stark 1988; Uchida and Isobe 1989; Terry and Whiting 2003; Chen et al. 2016, 2018; Wang et al. 2016, 2017a, 2019; Chen and Du 2017a, 2017b; Ding et al. 2019; Wipfler et al. 2019; South et al. 2020, 2021; Cao et al. 2021). In the superfamily Perloidea, the affinities of three families: Chloroperlidae, Perlidae and Perlodidae, have been proposed as a trichotomy (Nelson 1984; Zwick 2000), while Illies (1965) proposed Perlidae as sister to Chloroperlidae + Perlodidae. Most molecular data strongly supported the latter classification (Chen et al. 2016, 2018; Wang et al. 2016, 2017b, 2019; Chen and Du 2017b, 2017c; Ding et al. 2019; South et al. 2020, 2021; Cao et al. 2021).

At present, only three chloroperlid species, *Sweltsa longistyla* (Wu, 1938), *Suwallia bimaculata* (Okamoto, 1912) and *Suwallia errata* Li and Li, 2021, have been completely sequenced for mitogenomes (Chen and Du 2015b; Wang et al. 2018a; Cao et al. 2019a; Li et al. 2021). This limited mitogenomic data hinders the reconstruction of accurate phylogeny (Wu et al. 2014). In the present study, two complete mitogenomes from two genera, *Haploperla* and *Sweltsa*, were sequenced. Nucleotide composition, codon usage, RNA secondary structure, the evolutionary pattern among the PCGs, and structural elements within the control regions were analysed. Furthermore, a phylogenetic tree of the infraorder Systelognatha, excluding the family Kathroperlidae was also reconstructed based on the aligned PCGs using Bayesian inference (BI) and Maximum likelihood (ML) analyses.

2. Material and Methods

2.1. Specimen sample and DNA extraction

Adult specimens of *Haploperla japonica* were collected from Yoshino River, Minocho, Miyoshi City, Tokushima Prefecture, Japan, and adult samples of *Sweltsa* sp. were collected from a forested stream along the road No.328, Kasatori Mountain, Kumakogen, Ehime Prefecture, Japan. Species-level identification of *Sweltsa* species is unsure, due to taxonomic problems in Japanese represen-

tatives of the genus (Shimizu et al. 2005). All specimens were fixed in 100% ethanol and stored at -20°C. The examined specimens of *H. japonica* (No. VHem-0114) and *Sweltsa* sp. (No. VHem-0116) are deposited in the Entomological Museum of Henan Institute of Science and Technology (HIST), Henan Province, China. The full genomic DNA was isolated from thoracic muscle using the QIAamp DNA Blood Mini Kit (Qiagen, Germany) and maintained at -20°C prior to PCR analysis.

2.2. Genome sequencing, assembly and annotation

The full mitogenomes of both Chloroperlidae species were amplified and sequenced as described in previous studies (Li et al. 2017; Wang et al. 2017a, 2017b; Liu et al. 2018) and uploaded in GenBank with the accession numbers: OL351265 (*Haploperla japonica*) and OL351266 (*Sweltsa* sp.) (Table 1). The program BioEdit version 7.0.5.3 was used to assemble raw sequences into contigs (Hall 1999). tRNA genes were identified with the software ARWEN using default setting (Laslett and Canbäck 2008). PCG and two rRNA genes were identified using BLAST searches in NCBI (<http://www.ncbi.nlm.nih.gov>), and confirmed by alignments of homologous genes from previously published stonefly mitogenomes. MEGA 5.0 was used to calculate nucleotide composition and codon usage for PCGs (Tamura et al. 2011). Strand asymmetry was measured by the following formulas: AT skew= [A-T]/[A+T] and GC skew= [G-C]/[G+C] (Perna and Kocher 1995). Tandem repeats in the A+T-rich region were identified using Tandem Repeats Finder (<http://tandem.bu.edu/trf/trf.advanced.submit.html>).

2.3. Phylogenetic analysis

We used forty previously published and the two newly sequenced mitogenomes for phylogenetic analyses (Table 1). Two species from the family Nemouridae, *Amphinemura yao* Mo, Yang, Wang and Li, 2017 and *A. longispina* (Okamoto, 1922), were used as outgroup taxa (Table 1). Each PCG was aligned separately using the MAFFT algorithm implemented the TranslatorX online platform (Abascal et al. 2010). Poorly aligned regions were masked from the protein alignment before back-translation to nucleotides using GBlocks (implemented in the TranslatorX). Each rRNA gene was aligned separately by MAFFT v7.0 using G-INS-I strategy (Katoh and Standley 2013), and ambiguously aligned positions masked by GBlocks v0.91b (Castresana 2000).

We assembled the “PCG13 matrix” (11,049 bp in total) for the phylogenetic analyses, including 13 PCGs. GTR+I+G was the best-fit model for the nucleotide sequence alignments (jModelTest 0.1.1, Posada 2008). Bayesian analyses were conducted on the PCG13 dataset partitioned by gene was conducted Bayesian inference (BI), using MrBayes 3.2.6 and maximum likelihood (ML) using RAxML-HPC2 8.1.11 (Stamatakis 2006; Ronquist

Table 1. Classification of all Plecoptera species with sequenced mitogenomes, their genome lengths and accession numbers at Genbank.

Infraorder	Superfamily	Family	Subfamily	Species	Number (bp)	Accession Number	
Systellognatha	Perloidea	Chloroperlidae	Chloroperlinae	<i>Haploperla japonica</i>	16,012	OL351265	
				<i>Sweltsa</i> sp.	15,893	OL351266	
				<i>Sweltsa longistyla</i>	16,151	KM216826	
				<i>Suwallia errata</i>	16,146	MF198253	
				<i>Suwallia bimaculata</i>	16,125	MN121757	
		Perlidae	Acroneuriinae	<i>Acroneuria hainana</i>	15,804	NC_026104	
				<i>Acroneuria carolinensis</i>	15,718	MN969989	
				<i>Caroperla siveci</i>	15,353	MG677942	
				<i>Calineuria stigmatica</i>	15,070	MG677941*	
				<i>Flavoperla hatakeyamae</i>	15,730	MN821010	
Pteronarcyoidea	Pteronarcyoidea	Perlidae	Perlinae	<i>Flavoperla</i> sp. YW-2019	15,796	MN419916	
				<i>Flavoperla</i> sp. YZD-2020	15,805	MK905206*	
				<i>Niponiella limbatella</i>	15,924	MK686067	
				<i>Perlesta teaysia</i>	16,023	MN627432	
				<i>Sinacroneuria dabieshana</i>	15,752	MK492253	
				<i>Claassenia</i> sp.	15,774	MN419914	
				<i>Dinocras cephalotes</i>	15,666	NC_022843	
				<i>Etcocorema hochii</i>	15,854	MK905888	
				<i>Kamimuria chungnanshana</i>	15,943	NC_028076	
				<i>Kamimuria klapaleki</i>	16,077	MN400755	
Euholognatha	Nemouroidea (Outgroup)	Nemouridae	Isoperlinae	<i>Kamimuria wangi</i>	16,179	NC_024033	
				<i>Neoperlops gressitti</i>	15,699	MN400756	
			Perlodidae	<i>Paragnetina indentata</i>	15,885	MN627431	
				<i>Neoperla</i> sp.	15,667	KX091859*	
				<i>Neoperla ignacsiveci</i>	15,777	KX091858	
				<i>Togoperla limbata</i>	15,915	MN969990	
				<i>Togoperla</i> sp.	15,723	KM409708	
				<i>Isoperla bilineata</i>	15,048	MF716959	
				<i>Isoperla eximia</i>	16,034	MG910457	
				<i>Perlodes</i> sp.	16,039	MF197377	
			Peltoperlidae	<i>Pseudomegarcs japonica</i>	16,067	MG910458	
				<i>Cryptoperla stilifera</i>	15,633	KC952026*	
				<i>Peltoperlopsis cebuano</i>	15,790	MK387068	
				<i>Soliperla</i> sp.	15,877	MF716958	
			Pteronarcyidae	<i>Microperlinae</i>	<i>Microperla geei</i>	15,216	MN096323
				<i>Pteronarcys princeps</i>	16,004	NC_006133	
				<i>Pteronarcella badia</i>	15,585	NC_029248	
				<i>Styloperla</i> sp.	15,416	KR088971*	
			Styloperlidae	<i>Styloperla spinicercia</i>	16,129	KX845569	
				<i>Cerconychia flectospina</i>	15,188	MF100783*	
				<i>Amphinemura yao</i>	15,876	MH085447	
			Amphinemurinae	<i>Amphinemura longispina</i>	15,709	MH085446	

*Incomplete genome sequence

et al. 2012). For the BI analyses, two simultaneous runs of 10 million generations were conducted, sampling every 1,000 generations and a burn-in rate of 25%. Stationarity was assessed using Tracer v.1.5 (Estimated sample size > 200). Bootstrap ML analyses of 1,000 replicates was performed with the fast ML method conducted in RAxML using the GTRGAMMA model for nucleotide sequences.

3. Results and Discussion

3.1. General characters of mitogenomes and the base composition

In this study, the full mitogenomes of two stonefly species in the subfamily Chloroperlinae: *Haploperla japonica* and *Sweltsa* sp., were studied for the first time (Table 2, Fig. 1). The new complete mitogenomes were 15,893 (*Swelt-*

Table 2. The size of nucleotide composition of the *Haploperla japonica* and *Sweltsa* sp. mitogenomes.

Family	Subfamily	Species	Genome	PCGs	tRNAs	lrRNA	srRNA	Control region
Chloroperlidae	Chloroperlinae	<i>Haploperla japonica</i>	16012	11247	1483	1324	776	1118
		<i>Sweltsa</i> sp.	15893	11244	1488	1325	799	1011

Table 3. Mitogenome structures of *Haploperla japonica* and *Sweltsa* sp.

Gene	<i>Haploperla japonica</i>				<i>Sweltsa</i> sp.		
	Direction	Location	Size	IGN	Location	Size	IGN
tRNA ^{Ile}	F	1–67	67		1–67	67	
tRNA ^{Gln}	R	65–133	69	-3	65–133	69	-3
tRNA ^{Met}	F	139–207	69	5	136–204	69	2
ND2	F	208–1242	1035	0	205–1239	1035	0
tRNA ^{Trp}	F	1241–1308	68	-2	1238–1305	68	-2
tRNA ^{Cys}	R	1301–1367	67	-8	1298–1365	68	-8
tRNA ^{Tyr}	R	1368–1433	66	0	1366–1433	68	0
COI	F	1426–2982	1557	-8	1426–2982	1557	-8
tRNA ^{Leu(UUR)}	F	3003–3068	66	20	2995–3060	66	12
COII	F	3100–3783	688	31	3073–3760	688	12
tRNA ^{Lys}	F	3788–3858	71	0	3761–3831	71	0
tRNA ^{Asp}	F	3858–3926	69	-1	3831–3899	69	-1
ATP8	F	3927–4088	162	0	3900–4058	159	0
ATP6	F	4082–4759	678	-7	4052–4729	678	-7
COIII	F	4759–5547	789	-1	4729–5517	789	-1
tRNA ^{Gly}	F	5550–5615	66	2	5517–5582	66	2
ND3	F	5616–5969	354	0	5583–5936	354	0
tRNA ^{Ala}	F	5968–6033	66	-2	5935–6000	66	-2
tRNA ^{Arg}	F	6034–6099	66	0	6001–6066	66	0
tRNA ^{Asn}	F	6103–6168	66	3	6071–6136	66	4
tRNA ^{Ser(AGN)}	F	6169–6235	67	0	6137–6203	67	0
tRNA ^{Glu}	F	6236–6301	66	0	6204–6269	66	0
tRNA ^{Phe}	R	6316–6380	65	14	6275–6339	65	5
ND5	R	6381–8115	1735	0	6340–8074	1735	0
tRNA ^{His}	R	8116–8182	67	0	8075–8142	68	0
ND4	R	8187–9527	1341	4	8148–9488	1341	5
ND4L	R	9521–9817	297	-7	9482–9778	297	-7
tRNA ^{Thr}	F	9820–9886	67	2	9781–9848	68	2
tRNA ^{Pro}	R	9888–9955	68	1	9850–9917	68	1
ND6	F	9957–10481	525	1	9919–10443	525	1
CytB	F	10481–11617	1137	-1	10443–11579	1137	-1
tRNA ^{Ser(UCN)}	F	11616–11685	70	-2	11579–11648	70	-1
ND1	R	11706–12656	951	20	11670–12620	951	21
tRNA ^{Leu(CUN)}	R	12658–12723	66	1	12622–12687	66	1
lrRNA	R	12724–14047	1324	0	12688–14012	1325	0
tRNA ^{Val}	R	14048–14118	71	0	14013–14083	71	0
srRNA	R	14119–14894	776	0	14084–14882	799	0
A+T-rich region		14895–16012	1118	0	14883–15893	1011	0

sa sp.) and 16,012 bp (*H. japonica*) in length (Fig. 1, Table 2). Genome sizes of these mitogenomes were in the middle of the range compared to mitogenomes of other stoneflies, that range from 15,048 bp (*Isoperla bilineata*) (Chen et al. 2018) to 16,602 bp (*Nemoura nankinensis*) (Chen and Du 2017a). Control region variation is the primary factor influencing size variation in Plecoptera mitogenomes (Table 2). The typical gene content found in most insects were

found in these species: 37 mitochondrial genes (13PCGs, 22tRNAs and 2 rRNAs) and a large control region (Fig. 1). Twenty-three genes (9 PCGs and 14 tRNAs) were transcribed on the majority strand (J-strand), and the remaining fourteen genes (4 PCGs, 8 tRNAs, and 2 rRNAs) were encoded on the minority strand (N-strand) (Fig. 1).

Overlap between gene regions ranged from one to eight bp in length. Both species had seven bp overlaps




Figure 1. Map of the mitogenomes of *Haploperla japonica* and *Sweltsa* sp. Direction of gene transcription is indicated by the arrows. PCGs are shown as blue arrows, rRNA genes as purple arrows, tRNA genes as red arrows and CR as gray arrows. tRNA genes are labeled according to single-letter IUPAC-IUB abbreviations (L1: UUR, L2: CUN, S1: AGN, S2: UCN). The GC content is plotted using a black sliding window, as the deviation from the average GC content of the entire sequence. GC skew is plotted as the deviation from the average GC skew of the entire sequence.

in (ATP6–ATP8) and (ND4–ND4L), “ATGATAA” and “TTAACAT”, respectively. This phenomenon was also found in the mitogenome of many insects (Yang et al. 2017; Cao et al. 2019a). An intergenic spacer sequence was found to be located between *tRNA^{Leu(UUR)}* and *COII* in the two species, ranging from 12 bp in *Sweltsa* sp. to 31 bp in *Haploperla japonica* (Table 3).

3.2. Nucleotide composition

The A+T content in the *Haploperla japonica* and the *Sweltsa* sp. mitogenome is detailed in Table 4. The A+T content of *Sweltsa* sp. is similar to the base compositional biases reported for other chloroperlid species, that range from 66.7% (*Suwallia talalajensis* Zhiltzova, 1976) to 69.6% (*Sweltsa longistyla* (Wu, 1938)) (Chen and Du 2015b; Wang et al. 2018a), whereas A+T content in *H. japonica* is larger than other chloroperlids. The four major genome partitions (PCGs, tRNAs, rRNAs, and CR) and the whole mitogenome of both Chloroperlinae species were distinctly biased towards A and T. Among the four major partitions across Chloroperlinae species, PCGs consistently had the lowest A+T content, while the control region had the highest. A+T content of PCGs encoded on the minority strand (PCGs–N) was also higher than that of PCGs encoded on the majority strand (PCGs–J) (Table 4).

Both Chloroperlinae species showed positive AT-skew and negative GC-skew for the whole mitogenome. Total mitogenome of *Haploperla japonica*, the AT-skew and GC-skew were 0.065 and –0.223, respectively, whereas in *Sweltsa* sp., the AT-skew and GC-skew were 0.087

and –0.254, respectively. Slight T-skews and moderate C-skews were found in the majority strand PCGs while the reverse phenomenon was found for minority strand PCGs: both T-skews and G-skews were marked. For tRNAs and rRNAs, G-skews varied from slight to marked, while T-skews were slight to moderate. Modest A-skew and marked C-skew were found in the CR (Table 4).

For the majority strand PCGs, most known metazoan mitogenomes have positive AT-skew and negative GC-skew (Wei et al. 2010a). However, both Chloroperlinae species had negative AT-skews and negative GC-skews for the majority strand PCGs (Table 4). This difference has been noted for other published stonefly mitogenomes (Wu et al. 2014; Chen and Du 2017a, 2017c; Wang et al. 2017a, 2017b), as well as in other insects such as Philopteridae (bird lice), Aleyrodidae (whiteflies) and Braconidae (wasps); this phenomenon may be a result from replication direction charges (Wei et al. 2010a). For both Chloroperlinae species, the four major partitions (PCGs, tRNAs, rRNAs and CR) all had the same trends in the A+T content, AT and GC-skews which displayed similar patterns and were consistent with the common base composition biases of insect mitogenomes.

3.3. Protein-coding genes and codon usage

The total length of PCGs in the studied mitogenomes was 11,247 bp (*Haploperla japonica*) and 11,244 bp (*Sweltsa* sp.) (Table 2). In both mitogenomes from Chloroperlinae, typical start codons ATN (Met/Ile) were the most commonly used start codons. Additionally, TTG was inferred

Table 4. The nucleotide composition of the *Haploperla japonica* and *Sweltsa* sp. mitogenomes.

Region		<i>H. japonica</i>	<i>Sweltsa</i> sp.
Whole mitogenome	A+T%	69.7	68.2
	AT-skew	0.065	0.087
	GC-skew	-0.223	-0.254
PCGs	A+T%	68.2	66.5
	AT-skew	-0.173	-0.164
	GC-skew	-0.009	-0.047
PCGs-J	A+T%	66.4	64.5
	AT-skew	-0.096	-0.063
	GC-skew	-0.179	-0.235
PCGs-N	A+T%	71.0	69.6
	AT-skew	-0.289	-0.312
	GC-skew	0.307	0.305
tRNAs	A+T%	69.3	68.0
	AT-skew	-0.007	-0.022
	GC-skew	0.083	0.109
rRNAs	A+T%	72.4	72.1
	AT-skew	-0.100	-0.119
	GC-skew	0.283	0.318
CR	A+T%	80.6	80.0
	AT-skew	0.083	0.068
	GC-skew	-0.235	-0.257

Table 5. The start and stop codons of the *Haploperla japonica* and *Sweltsa* sp. mitogenomes.

Species	ND2		COI		COII		ATP8		ATP6		COIII		ND3	
	Start	Stop	Start	Stop	Start	Stop	Start	Stop	Start	Stop	Start	Stop	Start	Stop
<i>H. japonica</i>	ATG	TAA	ATT	TAA	ATG	T-	ATG	TAA	ATG	TAA	ATG	TAA	ATT	TAG
<i>Sweltsa</i> sp.	ATG	TAA	ATT	TAA	ATG	T-	ATG	TAA	ATG	TAA	ATG	TAA	ATC	TAG
Species	ND5		ND4		ND4L		ND6		CYTB		ND1			
	Start	Stop	Start	Stop	Start	Stop	Start	Start	Start	Stop	Start	Stop	Start	Stop
<i>H. japonica</i>	GTG	T-	ATG	TAA	ATG	TAA	ATT	TAA	ATG	TAG	TTG	TAG		
<i>Sweltsa</i> sp.	GTG	T-	ATG	TAA	ATG	TAA	ATC	TAA	ATG	TAA	TTG	TAG		

as a start codon for *ND1*, and GTG as the start codon for *ND5* in both species. This phenomenon has been found in other stoneflies, and TTG is used in other insect species (Bae et al. 2004; James and Andrew 2006; Sheffield et al. 2008; Wei et al. 2010b; Elbrecht and Leese 2015; Elbrecht et al. 2015; Huang et al. 2015; Sproul et al. 2015; Chen et al. 2016; Chen and Du 2017c; Wang et al. 2017a, 2017b; Cao et al. 2019b). In both species, *ATP6*, *ATP8*, *COII*, *COIII*, *CYTB*, *ND2*, *ND4* and *ND4L* started with ATG (Met), while *COI*, *ND3* and *ND6* started with ATT (Ile) (Table 5).

The most commonly used stop codon in both Chloroperlinae species was TAA, found in *ATP6*, *ATP8*, *COI*, *COIII*, *ND2*, *ND4*, *ND4L* and *ND6*. In both species, the stop codon TAG was used by *CYTB*, *ND1* and *ND3* (except *CYTB* of *Sweltsa* sp., which used TAA). *COII* and *ND5* used a partial stop codon T in both species (Table 5). This phenomenon has also been recorded from other published stoneflies (James and Andrew 2006; Qian et al. 2014; Elbrecht and Leese 2015; Elbrecht et al. 2015; Huang et al. 2015; Wu et al. 2014; Chen et al. 2016; Chen and Du 2017c; Wang et al. 2017a, 2017b; Cao et al.

2019b). Truncated stop codons occur in many arthropod mitogenomes, and it is corrected by post-transcriptional polyadenylation (Ojala et al. 1981; Wolstenholme 1992). The two species had the same start and stop codons in ten genes (*COI*, *COII*, *COIII*, *ATP8*, *ATP6*, *ND1*, *ND2*, *ND4*, *ND4L* and *ND5*), while the remaining three genes (*ND3*, *ND6* and *CYTB*) only shared either the same start or stop codon (Table 5).

The genome-wide AT bias is reflected in codon usage. RSCU (relative synonymous codon usage) was calculated to identify the predominant synonymous codon (Grantham et al. 1980; Meganathan et al. 2012) (Fig. 2). The high AT base compositional biases was shown in the RSCU statistics for both Chloroperlinae mitogenomes. Codons ending with an A or U were favoured both four-fold and two-fold degenerate codons. RSCU demonstrates that codon usage patterns of these two stonefly species was consistent with other published Plecoptera species. All most commonly used codons: AAU (Asn), AUU (Ile), UUA (Leu1), AAA (Lys), AUA (Met), UUU (Phe) and UAU (Tyr), were always biased towards A and T at third codon positions (Fig. 2, Table 5).




Figure 2. Relative synonymous codon usage (RSCU) in *Haploperla japonica* and *Sweltsa* sp. mitogenomes. Codon families are provided on the X-axis.

3.4. Transfer and ribosomal RNAs

In the present study, the tRNA genes in both species ranged from 65 to 71 bp, and comprised 9.26% (1,483 bp in *H. japonica*) and 9.36% (1,488 bp in *Sweltsa* sp.) of the complete mitogenomes (Tables 2 and 3). *tRNA^{Ser(AGN)}*, which could not be detected by ARWEN software, was identified by alignment with homologous genes from other published stoneflies (James and Andrew 2006; Chen et al. 2016; Wang et al. 2017a, 2018a, 2018b; Cao et al. 2019b; Guo et al. 2021). Most tRNAs had the typical cloverleaf structure, except *tRNA^{Ser(AGN)}* that lacks the dihydrouridine (DHU) arm that formed a simple loop. In most insect mitogenomes, it is common for *tRNA^{Ser(AGN)}* to lack the DHU stem (Cameron and Whiting 2007; Wan et al. 2013). The length of the different nucleotides of tRNAs between *H. japonica* and *Sweltsa* sp. ranged from 1 bp (*tRNA^{Ser(AGN)}*, *tRNA^{Ser(UCN)}*, *tRNA^{Val}*) to 13 bp (*tRNA^{His}*). By comparison, the *tRNA^{Cys}* and *tRNA^{His}* had the most variation with more than ten indels or substitutions; the nucleotide insertion-deletion appears in *tRNA^{Cys}*, *tRNA^{His}*, *tRNA^{Phe}*, *tRNA^{Thr}*, *tRNA^{Tyr}* (Fig. 3).

Based on the secondary structures of tRNA, eight mismatched base pairs are found in *H. japonica*, which are U-U (2 bp), A-A (3 bp) and A-C (3 bp) (Fig. 3). And they have 24 G-U pairs, 7 located in the AA, 7 in the DHU, 8 in the AC and 2 in the TYC stem. In *Sweltsa* sp., there are nine mismatches in tRNAs: U-U (1 bp), A-A (5 bp), C-C (1 bp) and A-C (2 bp) (Fig. 3); the 29 G-U pairs are located in the AA (5 bp), the DHU (8 bp), the AC (12 bp) and the TYC stem (4 bp).

Similar to the mitogenomes of most other insects, the large and small rRNA subunits (*lrRNA* and *srRNA*) in both species were located between *tRNA^{Leu(CUN)}* – *tRNA^{Val}* (*lrRNA*) and *tRNA^{Val}* – the control region (*srRNA*) (Fig. 1, Table 3). The lengths of *lrRNA* and *srRNA* of *Haploperla japonica* were 1324 bp and 776 bp, and the lengths of *lrRNA* and *srRNA* of *Sweltsa* sp. were 1325 bp and 799 bp (Table 2). Secondary structures of both *lrRNA* and *srRNA* were highly similar between these species (Pairwise sequence identity was 87.82% (*lrRNA*) and 87.98% (*srRNA*)).

Haploperla japonica was used as model for comparison of rRNA secondary structures. *lrRNA* consists of 5 structural domains (I–II, IV–VI), while domain III was absent, as usual in arthropods (Fig. 4) (Cannone et al. 2002). Domains I, II and IV were variable between these species, while eight helices (H2064, H2347, H2395, H2455, H2507, H2520, H2547 and H2588) within domain V had high similarity (Wang et al. 2017a).

srRNA consists of three domains (Fig. 5). Domains I and II were more variable than domain III in both species, while Helix 1399 region was the most conserved (Wang et al. 2017a). Furthermore, *Sweltsa* sp. possessed 21 more bp nucleotides in Helix 577 than *Haploperla japonica*.

3.5. The control region

In a previous study, the A+T-rich region (CR) was found to contain essential elements for the initiation of tran-




Figure 3. Secondary structures of 22 tRNAs of *Haploperla japonica* and *Sweltsa* sp. All tRNAs are labeled with the abbreviations of their corresponding amino acids. Dashes (—) indicate Watson-Crick base pairing and dots (•) indicate G-U base pairing. Both species are represented in gray; blue circle only on *H. japonica*, yellow only on *Sweltsa* sp.; green means not on *H. japonica*, red means not on *Sweltsa* sp.

scription and for replication (Zhang and Hewitt 1997). *Haploperla japonica* mitogenome had an A+T-rich region of 1,118 bp, with an A+T content of 80.6%; while *Sweltsa* sp. it was 1,011 bp long, with an A+T content of 80.0% (Tables 2 and 4). The A+T-rich region of both spe-

cies is located between the *srRNA* and *tRNA^{Leu}* gene cluster (Fig. 1). The following structural elements were found in the control region of *H. japonica* and *Sweltsa* sp. (Fig. 6): (1) a leading sequence (588 bp (*H. japonica*), 496 bp (*Sweltsa* sp.)) adjacent to *srRNA*; (2) two large tandem




Figure 4. Predicted secondary structure of the *18S rRNA* in *Haploperla japonica* and *Sweltsa* sp. Roman numerals represent the conserved domain structures. Dashes (–) indicate Watson-Crick base pairings and dots (•) indicate G-U base pairing. Different nucleotides of the two species are shown in red. Missing nucleotides in *H. japonica* are indicated by red arrows, missing nucleotides in *Sweltsa* sp. are shown in blue.

repeats (TRs), while (3) the remaining control region was 321 bp (*H. japonica*) and 372 bp (*Sweltsa* sp.).

3.6. Phylogenetic relationships

Phylogenetic relationships were reconstructed for the clade Systellognatha using 42 species: 36 complete (including the two species herein studied) and 6 nearly complete mitogenome sequences (Table 1). *Amphinemura yao* Mo, Yang, Wang and Li, 2017 and *A. longispina* (Okamoto, 1922) were used as outgroup taxa. Bayesian inference (BI) and maximum likelihood (ML) analyses, of the aligned 13 PCGs dataset, generated trees with identical topologies (Fig. 7).

Systellognatha includes seven families: superfamily Perloidea with Chloroperlidae, Kathroperlidae, Perlidae, Perlodidae; and superfamily Pteronarcyoidea with Peltoperlidae, Pteronarcyidae, Styloperlidae; all except Kathroperlidae (no mitogenomes yet sequenced) were represented in our study. We obtained all families as monophyletic (colors in Fig. 7). Our analyses strongly supported Peltoperlidae as the sister group of a clade containing the five other families. Similar relationships were recovered in previous phylogenetic analyses (Chen et al. 2018; South et al. 2020, 2021; Cao et al. 2021). A sister-group relationship between Perloidea (monophyletic herein: Fig. 7) and Pteronarcyoidea was not supported, as the latter was obtained as paraphyletic, which is in-

consistent with morphological hypotheses (Zwick 2000). Phylogenetic analyses based on morphological characters proposed a clade Peltoperlidae + Styloperlidae, with Pteronarcyidae as their sister group (Zwick 2000). According to the present studies, Styloperlidae is sister to Pteronarcyidae rather than Peltoperlidae albeit with poor nodal support (as in Ding et al. 2019; Wang et al. 2019; Cao et al. 2019a; 2021). Perlidae was sister to Chloroperlidae + Perlodidae with high support (Fig. 7) (confirming Ricker 1952; Illies 1965; Chen et al. 2018; Wang et al. 2019, 2018c; South et al. 2020, 2021). Our analyses found relationships within Systellognatha to be (((Chloroperlidae + Perlodidae) + Perlidae) + (Styloperlidae + Pteronarcyidae)) + Peltoperlidae (Fig. 7). Recent phylogenetic analyses based on mitochondrial data have inferred the same topology (Chen et al. 2018; Cao et al. 2021), however, our results differ from the older morphology-based classification. The different results may be due to the limited mitogenomic data yet published about the Systellognatha, being especially scarce in Pteronarcyidae and Styloperlidae, and lacking in Kathroperlidae.

At the generic level, one pattern is the disjunct phylogenetic positions for Asian and North American members of the same genus (Fig. 7). For example, regarding *Isoperla* Banks, 1906, *Pseudomegarctys japonica* Kohno, 1946 was sister to the Asian *I. eximia* Zapekina-Dulkeit, 1975, rather than the North American *I. bilineata* (Say, 1823), the type species of genus *Isoperla*. Additionally, regarding *Acroneuria* Pictet, 1841, *Sinacroneuria dabieshana*




Figure 5. Predicted secondary structure of the *srRNA* in *Haploperla japonica* and *Sweltsa* sp. Roman numerals denote the conserved domain structure. Dashes (—) indicate Watson-Crick base pairing and dots (•) indicate G-U base pairing. Different nucleotides of the two species are shown in red. Missing nucleotides in *H. japonica* are indicated by red arrows, missing nucleotides in *Sweltsa* sp. are shown in blue.




Figure 6. Structure elements found in the control region of *Haploperla japonica* and *Sweltsa* sp.




Figure 7. Phylogenetic tree inferred from the sequences of the mitogenomes of 40 Systelognatha species and 2 outgroup species. Numbers at the nodes are Bayesian posterior probabilities (left) and ML bootstrap values (right).

Li and Murányi, 2014 was sister to the Asian species *A. hainana* Wu, 1938, rather than the North American *A. carolinensis* (Banks, 1905). Regarding their morphology, both *Pseudomegarctys* and *Sinacroneuria* are rather distinctive genera, in addition, *Pseudomegarctys* belongs to the subfamily Perlodinae Klapálek, 1909, while *Isoperla* belongs to Isoperlinea Frison, 1942. Inclusion of these genera in *Isoperla* and *Acroneuria*, respectively, is not likely, and additional molecular studies are needed to clarify their phylogenetic position.

4. Competing Interests

The authors have declared that no conflict of interest exists.

5. Acknowledgments

The research was partially supported by the National Natural Science Foundation of China (No. 31970402), the Graduate Education Innovation Training Base Project of Henan Province in 2021 (107020221005) and the Program for Science & Technology Innovation Talents in Universities of Henan Province (No. 21HASTIT042).

6. References

- Abascal F, Zardoya R, Telford MJ (2010) TranslatorX: multiple alignment of nucleotide sequences guided by amino acid translations. *Nucleic Acids Research* 38: 7–13. <https://doi.org/10.1093/nar/gkq291>
- Alexander KD, Stewart KW (1999) Revision of the genus *Suwallia* Ricker (Plecoptera: Chloroperlidae). *Transactions of the American Entomological Society* 125: 185–250.

Bae JS, Kim I, Sohn HD, Jin BR (2004) The mitochondrial genome of the firefly, *Pyrocoelia rufa*: complete DNA sequence, genome organization, and phylogenetic analysis with other insects. *Molecular Phylogenetics and Evolution* 32(3): 978–985. <https://doi.org/10.1016/j.ympev.2004.03.009>

Boore JL (1999) Animal mitochondrial genomes. *Nucleic Acids Research* 27: 1767–1780. <https://doi.org/10.1093/nar/27.8.1767>

Cameron SL, Whiting MF (2007) Mitochondrial genomic comparisons of the subterranean termites from the genus *Reticulitermes* (Insecta: Isoptera: Rhinotermitidae). *Genome* 50: 188–202. <https://doi.org/10.1139/G06-148>

Cannone JJ, Subramanian S, Schnare MN, Collett JR, D'Souza LM, Du Y, Feng B, Lin N, Madabusi LV, Müller KM, Pande N, Shang Z, Yu N, Gutell RR (2002) The comparative RNA web (CRW) site: An online database of comparative sequence and structure information for ribosomal, intron, and other RNAs. *BMC Bioinformatics* 3: 15.

Cao JJ, Wang Y, Chen MD, Yuan MY, Li WH (2019a) The complete mitochondrial genome of a stonefly species, *Suwallia bimaculata* (Plecoptera: Chloroperlidae). *Mitochondrial DNA Part B* 4(2): 2828–2829. <https://doi.org/10.1080/23802359.2019.1660263>

Cao JJ, Wang Y, Guo X, Wang GQ, Li WH, Murányi D (2021) Two complete mitochondrial genomes from Leuctridae (Plecoptera: Nemouroidea): Implications for the phylogenetic relationships among stoneflies. *Journal of Insect Science* 21(1): 16; 1–6. <https://doi.org/10.1093/jisesa/ieab009>

Cao JJ, Wang Y, Li WH (2019b) Comparative mitogenomic analysis of species in the subfamily Amphinemurinae (Plecoptera: Nemouridae) reveal conserved mitochondrial genome organization. *International Journal of Biological Macromolecules* 138: 292–301. <https://doi.org/10.1016/j.ijbiomac.2019.07.087>

Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17(4): 540–552. <https://doi.org/10.1093/oxfordjournals.molbev.a026334>

- Chen ZT, Du YZ (2015a) A new species of *Suwallia* (Plecoptera: Chloroperlidae) from China. *Zootaxa* 4018(2): 297–300. <https://doi.org/10.11646/zootaxa.4018.2.9>
- Chen ZT, Du YZ (2015b) Comparison of the complete mitochondrial genome of the stonefly *Sweltsa longistyla* (Plecoptera: Chloroperlidae) with mitogenomes of three other stoneflies. *Gene* 558(1): 82–87. <https://doi.org/10.1016/j.gene.2014.12.049>
- Chen ZT, Du YZ (2017a) First mitochondrial genome from Nemouridae (Plecoptera) reveals novel features of the elongated control region and phylogenetic implications. *International Journal of Molecular Sciences* 18(5): 996. <https://doi.org/10.3390/ijms18050996>
- Chen ZT, Du YZ (2017b) The first two mitochondrial genomes from Taeniopterygidae (Insecta: Plecoptera): Structural features and phylogenetic implications. *International Journal of Biological Macromolecules* 111: 70–76. <https://doi.org/10.1016/j.ijbiomac.2017.12.150>
- Chen ZT, Du YZ (2017c) Complete mitochondrial genome of *Capnia zjinshana* (Plecoptera: Capniidae) and phylogenetic analysis among stoneflies. *Journal of Asia-Pacific Entomology* 20(2): 305–312. <https://doi.org/10.1016/j.aspen.2017.01.013>
- Chen ZT, Wu HY, Du YZ (2016) The nearly complete mitochondrial genome of a stonefly species, *Styloperla* sp. (Plecoptera: Styloperlidae). *Mitochondrial DNA Part A* 27: 2728–2729. <https://doi.org/10.3109/19401736.2015.1046166>
- Chen ZT, Zhao MY, Xu C, Du YZ (2018) Molecular phylogeny of Systellognatha (Plecoptera: Arctoperlaria) inferred from mitochondrial genome sequences. *International Journal of Biological Macromolecules* 111: 542–547. <https://doi.org/10.1016/j.ijbiomac.2018.01.065>
- DeWalt RE, Maehr MD, Hopkins H, Neu-Becker U, Stueber G (2021) Plecoptera Species File Online. Version 5.0/5.0. Available from: <http://Plecoptera.SpeciesFile.org> (accessed 13 August 2021)
- Ding SM, Li WH, Wang Y, Cameron SL, Murányi D, Yang D (2019) The phylogeny and evolutionary timescale of stoneflies (Insecta: Plecoptera) inferred from mitochondrial genomes. *Molecular Phylogenetics and Evolution* 135: 123–135. <https://doi.org/10.1016/j.ympev.2019.03.005>
- Elbrecht V, Leese F (2015) The mitochondrial genome of the Arizona Snowfly *Mesocapnia arizonensis* (Plecoptera, Capniidae). *Mitochondrial DNA* 27(5): 1–2. <https://doi.org/10.3109/19401736.2015.1018223>
- Elbrecht V, Poettker L, John U, Leese F (2015) The complete mitochondrial genome of the stonefly *Dinocras cephalotes* (Plecoptera, Perlidae). *Mitochondrial DNA* 26(3): 469–470. <https://doi.org/10.3109/19401736.2013.830301>
- Frison TH (1942) Studies of North American Plecoptera, with special reference to the fauna of Illinois. *Bulletin of the Illinois Natural History Survey* 22: 235–355.
- Gissi C, Iannelli F, Pesole G (2008) Evolution of the mitochondrial genome of Metazoa as exemplified by comparison of congeneric species. *Heredity* 101: 301–320. <https://doi.org/10.1038/hdy.2008.62>
- Guo X, Dong XJ, Yang M, Gao S, Wang Y (2021) Mitochondrial genome of *Arcynopteryx dichroa* (McLachlan, 1872) (Plecoptera: Perlodidae) and phylogenetic analysis. *Mitochondrial DNA Part B – Resources* 6: 2433–2435. <https://doi.org/10.1080/23802359.2021.1955764>
- Grantham R, Gautier C, Gouy M (1980) Codon catalog usage and the genome hypothesis. *Nucleic Acids Research* 8: r49–r62.
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Huang MC, Wang YY, Liu XY, Li WH, Kang ZH, Wang K, Li XK, Yang D (2015) The complete mitochondrial genome and its remarkable secondary structure for a stonefly *Acroneuria hainana* Wu (Insecta: Plecoptera, Perlidae). *Gene* 557(1): 52–60. <https://doi.org/10.1016/j.gene.2014.12.009>
- Illies J (1965) Phylogeny and zoogeography of the Plecoptera. *Annual Review of Entomology* 10: 117–140.
- James BS, Andrew TB (2006) Insect mitochondrial genomics 2: the complete mitochondrial genome sequence of a giant stonefly, *Pteronarcys princeps*, asymmetric directional mutation bias, and conserved plecopteran A + T-region elements. *Genome* 49: 815–824. <https://doi.org/10.1139/g06-037>
- Judson SW, Nelson CR (2012) A guide to Mongolian stoneflies (Insecta: Plecoptera). *Zootaxa* 3541: 1–118. <https://doi.org/10.11646/zootaxa.3541.1.1>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Klapálek F (1909) Plecoptera, Steinfliegen. Die Süßwasserfauna Deutschlands 8: 33–95.
- Laslett D, Canbäck B (2008) ARWEN: a program to detect tRNA genes in metazoan mitochondrial nucleotide sequences. *Bioinformatics* 24(2): 172–175. <https://doi.org/10.1093/bioinformatics/btm573>
- Li H, Leavengood Jr JM, Chapman EG, Burkhardt D, Song F, Jiang P, Liu J, Zhou XG, Cai WZ (2017) Mitochondrial phylogenomics of Hemiptera reveals adaptive innovations driving the diversification of true bugs. *Proceedings of the Royal Society B: Biological Sciences* 284: 1862. <https://doi.org/10.1098/rspb.2017.1223>
- Li WH, Murányi D, Li S (2015a) New species records of *Suwallia* Ricker, 1943 (Plecoptera: Chloroperlidae) from China, with description of the nymph of *S. decolorata* Zhiltzova and Levanidova, 1978. *Zootaxa* 3994(4): 556–564. <https://doi.org/10.11646/zootaxa.3994.4.4>
- Li WH, Murányi D, Li S (2015b) The first record of genus *Suwallia* Ricker, 1943 (Plecoptera: Chloroperlidae) from China. *Illiesia* 11(3): 23–28.
- Li WL, Wang YY, Wang Y, Li WH (2021) A new species of *Suwallia* Ricker, 1943 from Japan, and the identity of *Alloperla telekujensis* Šámal, 1939 (Plecoptera: Chloroperlidae). *Zootaxa* 5040(4): 575–581. <https://doi.org/10.11646/zootaxa.5040.4.7>
- Liu YQ, Song F, Jiang P, Wilson JJ, Cai WZ, Li H (2018) Compositional heterogeneity in true bug mitochondrial phylogenomics. *Molecular Phylogenetics and Evolution* 118: 135–144. <https://doi.org/10.1016/j.ympev.2017.09.025>
- Meganathan PR, Pagan HJT, McCulloch ES, Stevens RD, Ray DA (2012) Complete mitochondrial genome sequences of three bats species and whole genome mitochondrial analyses reveal patterns of codon bias and lend support to a basal split in Chiroptera. *Gene* 492: 121–129. <https://doi.org/10.1016/j.gene.2011.10.038>
- Nelson CH (1984) Numerical cladistic-analysis of phylogenetic relationships in Plecoptera. *Annals of the Entomological Society of America* 77: 466–473.
- Ojala D, Montoya J, Attardi G (1981) tRNA punctuation model of RNA processing in human mitochondria. *Nature* 290: 470–474. <https://doi.org/10.1038/290470a0>
- Perna NT, Kocher TD (1995) Patterns of nucleotide composition at four-fold degenerate sites of animal mitochondrial genomes. *Journal of Molecular Evolution* 41: 353–358.

- Posada D (2008) jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* 25(7): 1253–1256. <https://doi.org/10.1093/molbev/msn083>
- Qian YH, Wu HY, Ji XY, Yu WW, Du YZ (2014) Mitochondrial genome of the stonefly *Kamimuria wangi* (Plecoptera: Perlidae) and phylogenetic position of Plecoptera based on mitogenomes. *PLoS ONE* 9: e86328. <https://doi.org/10.1371/journal.pone.0086328>
- Ricker WE (1952) Systematic studies in Plecoptera. *Indiana University Publications Science Series* 18: 1–200.
- Ronquist F, Teslenko M, van der Mark PVD, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Sheffield NC, Song H, Cameron SL, Whiting MF (2008) A comparative analysis of mitochondrial genomes in Coleoptera (Arthropoda: Insecta) and genome descriptions of six new beetles. *Molecular Biology and Evolution* 25(11): 2499–2509. <https://doi.org/10.1093/molbev/msn198>
- Shimizu T, Inada K, Uchida S (2005) Plecoptera, in: Kawai T, Tanida K (Eds), *Aquatic Insects of Japan: Manual with keys and illustrations*. Tokai University Press, Kanagawa, pp. 237–290.
- South EJ, Skinner RK, DeWalt RE, Davis MA, Johnson KP, Teslenko AV, Lee JJ, Malison RL, Hwang YM, Bae YJ, Myers LW (2021) A new family of stoneflies (Insecta: Plecoptera), Kathroperlidae, fam. n., with a phylogenomic analysis of the Paraperlinae (Plecoptera: Chloroperlidae). *Insect Systematics and Diversity* 5(4): 1–27. <https://doi.org/10.1093/isd/ixab014>
- South EJ, Skinner RK, DeWalt RE, Kondratieff BC, Johnson KP, Davis MA, Lee JJ, Durfee RS (2020) Phylogenomics of the North American Plecoptera. *Systematic Entomology* 1–19. <https://doi.org/10.1111/syen.12462>
- Sproul JS, Houston DD, Nelson CR, Evans P, Crandall KA, Shiozawa DK (2015) Climate oscillations, glacial refugia, and dispersal ability: factors influencing the genetic structure of the least salmonfly, *Pteronarcella badia* (Plecoptera), in Western North America. *BMC Evolutionary Biology* 15: 1–18. <https://doi.org/10.1186/s12862-015-0553-4>
- Stamatakis A (2006) RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688–2690. <https://doi.org/10.1093/bioinformatics/btl446>
- Stewart KW, Stark BP (1988) Nymphs of North American stonefly genera (Plecoptera). Thomas Say Foundation 12: 1–436.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28(10): 2731–2739. <https://doi.org/10.1093/molbev/msr121>
- Terry MD, Whiting MF (2003) Phylogeny of Plecoptera: molecular evidence and evolutionary trends. *Entomologische Abhandlungen* 61: 130–131.
- Tracer version 1.5. <http://tree.bio.ed.ac.uk/software/tracer> (accessed 16 Feb 2018).
- Uchida S, Isobe Y (1989) Styloperlidae, stat. nov. and Microperlinae, subfam. nov. with a revised system of the family group Systellognatha (Plecoptera). *Spixiana* 12: 145–182.
- Wan X, Kim MI, Kim MJ, Kim I (2013) Complete mitochondrial genome of the free-living earwig, *Challia fletcheri* (Dermaptera: Pygidicranidae) and phylogeny of Polyneoptera. *PLoS One* 7: e42056. <https://doi.org/10.1371/journal.pone.0042056>
- Wang Y, Cao JJ, Li WH (2017a) The complete mitochondrial genome of the styloperlid stonefly species *Styloperlula spinicercia* Wu (Insecta: Plecoptera) with family-level phylogenetic analyses of the Pteronarcyoidea. *Zootaxa* 4243(1): 125–138. <https://doi.org/10.11646/zootaxa.4243.1.5>
- Wang Y, Cao JJ, Li WH (2018a) Complete mitochondrial genome of *Sutwallia teleckojensis* (Plecoptera: Chloroperlidae) and implications for the higher phylogeny of stoneflies. *International Journal of Molecular Science* 19(3): 680. <https://doi.org/10.3390/ijms19030680>
- Wang Y, Cao JJ, Lei BB, Li WH (2018b) The mitochondrial genome of a stonefly species, *Cerconychia sapa* (Plecoptera: Styloperlidae). *Conservation Genetics Resources* 10: 145–148. <https://doi.org/10.1007/s12686-017-0781-6>
- Wang Y, Cao JJ, Li N, Ma GY, Li WH (2019) The first mitochondrial genome from Scopuridae (Insecta: Plecoptera) reveals structural features and phylogenetic implications. *International Journal of Biological Macromolecules* 122: 893–902. <https://doi.org/10.1016/j.ijbiomac.2018.11.019>
- Wang Y, Cao JJ, Li WH, Chen XL (2017b) The mitochondrial genome of *Mesocapina daxingana* (Plecoptera: Capniidae). *Conservation Genetics Resources* 9: 639–642. <https://doi.org/10.1007/s12686-017-0745-x>
- Wang Y, Cao JJ, Murányi D, Li WH (2018c) Comparison of two complete mitochondrial genomes from Perlodidae (Plecoptera: Perlodea) and the family-level phylogenetic implications of Perlodea. *Gene* 675: 254–264. <https://doi.org/10.1016/j.gene.2018.06.093>
- Wang K, Ding SM, Yang D (2016) The complete mitochondrial genome of a stonefly species, *Kamimuria chungnanshana* Wu, 1948 (Plecoptera: Perlidae). *Mitochondrial DNA Part A* 27: 3810–3811. <https://doi.org/10.3109/19401736.2015.1082088>
- Wei S, Shi M, Chen X, Sharkey MJ, van Achterberg C, Ye G, He J (2010a) New views on strand asymmetry in insect mitochondrial genomes. *PLoS ONE* 5: e12708. <https://doi.org/10.1371/journal.pone.0012708>
- Wei SJ, Tang P, Zheng LH, Shi M, Chen XX (2010b) The complete mitochondrial genome of *Evania appendigaster* (Hymenoptera: Evaniidae) has low A+T content and a long intergenic spacer between atp8 and atp6. *Molecular Biology Reports* 37: 1931–1942. <https://doi.org/10.1007/s11033-009-9640-1>
- Wipfler B, Letsch H, Frandsen PB, Kapli P, Mayer C, Bartel D, Buckley TR, Donath A, Edgerly-Rooks JS, Fujita M, Liu SL, Machida R, Mashimo Y, Misof B, Niehuis O, Peters RS, Petersen M, Podsiadlowski L, Schütte K, Shimizu S, Uchifune T, Wilbrandt J, Yan E, Zhou X, Simon S (2019) Evolutionary history of Polyneoptera and its implications for our understanding of early winged insects. *PNAS* 116(8): 3024–3029. <https://doi.org/10.1073/pnas.1817794116>
- Wolstenholme DR (1992) Animal mitochondrial DNA: structure and evolution. *International Review of Cytology* 141: 173–216. [https://doi.org/10.1016/S0074-7696\(08\)62066-5](https://doi.org/10.1016/S0074-7696(08)62066-5)
- Wu HY, Ji XY, Yu WW, Du YZ (2014) Complete mitochondrial genome of the stonefly *Cryptoperla stilifera* Sivec (Plecoptera: Peltoperlidae) and the phylogeny of polyneopteran insects. *Gene* 537(2): 177–183. <https://doi.org/10.1016/j.gene.2013.12.044>
- Yang F, Jiang YL, Yang D, Liu XY (2017) Mitochondrial genomes of two Australian fishflies with an evolutionary timescale of Chauliodinae. *Scientific Reports* 7: 4481, 1–13. <https://doi.org/10.1038/s41598-017-04799-y>

- Zhang DX, Hewitt GM (1997) Insect mitochondrial control region: a review of its structure, evolution and usefulness in evolutionary studies. *Biochemical Systematics and Ecology* 25: 99–120. [https://doi.org/10.1016/S0305-1978\(96\)00042-7](https://doi.org/10.1016/S0305-1978(96)00042-7)
- Zwick P (2000) Phylogenetic system and zoogeography of the Plecop-
tera. *Annual Review of Entomology* 45: 709–746.