



Dasyophthalma (Lepidoptera: Nymphalidae: Satyrinae): systematics, distribution, and conservation perspectives of a butterfly genus endemic from the Brazilian Atlantic Forest

Augusto Henrique Batista Rosa¹, Eduardo de Proença Barbosa¹, Patrícia Avelino Machado¹
Ricardo Russo Siewert¹, André Victor Lucci Freitas¹

¹ Depto de Biologia Animal and Museu de Diversidade Biológica, Instituto de Biologia, Universidade Estadual de Campinas, Rua Bertrand Russell, s/n, CEP: 1202-1292, Campinas, São Paulo, Brazil

<https://zoobank.org/49DC3DD6-8D98-4399-8E2F-9F821F31833C>

Corresponding authors: André V. L. Freitas (baku@unicamp.br), Augusto H. B. Rosa (augustohbrosa@hotmail.com)

Received 17 October 2022

Accepted 30 March 2023

Published 23 May 2023

Academic Editors Ângelo Pinto, Martin Wiemers

Citation: Rosa AHB, Barbosa EP, Machado PA, Siewert RR, Freitas AVL (2023) *Dasyophthalma* (Lepidoptera: Nymphalidae: Satyrinae): systematics, distribution, and conservation perspectives of a butterfly genus endemic from the Brazilian Atlantic Forest. Arthropod Systematics & Phylogeny 81: 455–473. <https://doi.org/10.3897/asp.81.e96397>

Abstract

Dasyophthalma includes five species of medium-sized butterflies, all endemic from the Atlantic Forest of Brazil. All known species are univoltine and are dayflying, differently from other Brassolini that are mostly crepuscular. In despite of recent advances little is known about their natural history. Three out of the five species are included in the Brazilian Red List of threatened fauna and are of conservation concern. The present study provides for the first time a phylogenetic assessment of all *Dasyophthalma* species based on a molecular approach based on three loci. Also, the taxonomic status of *D. rusina delanira* was revised based on molecular data. In addition, up-to-date distributional data and conservation aspects of the threatened species from the genus are presented and discussed. The molecular phylogenetic analysis supports the monophyly of *Dasyophthalma*, with *Dynastor darius* as its sister-group, and, combined with a genetic divergence analysis, supported *Dasyophthalma delanira* **stat. rest.** as a valid name to species-level, sister-group to *D. geraensis* (and not a subspecies of *D. rusina*). The geographical range (extent of occurrence and area of occupancy) for all five species are presented, showing that these are very restricted for *D. delanira* **stat. rest.** and *D. geraensis*, following the distributions of the high-altitude forests. As much biological information about the genus is lacking, the present study can serve as a starting point for future studies on *Dasyophthalma*, adding information that can be crucial for future conservation actions and essential to assure the future of the threatened species in this genus.

Keywords

Brassolini, fruit-feeding butterflies, Neotropical, phylogeny, endangered species.

1. Introduction

The tribe Brassolini (Nymphalidae: Satyrinae) is a clade of exclusively Neotropical butterflies that currently includes 108 species distributed in 17 genera (Penz 2007; Shirai et al. 2016). Brassolini butterflies are notorious by

being large sized, including some of the largest Neotropical butterflies, and by presenting several species with crepuscular behavior, flying at dawn or in the first hours of the day (DeVries 1987; Freitas et al. 1997; Penz 2007).

The genus *Dasyophthalma* Westwood, 1851 includes species of conspicuous medium sized butterflies, all endemic from the Brazilian Atlantic Forest. All known species are univoltine, with adults usually flying during the summer season and unlike most brassolines, which have crepuscular habits, all species of *Dasyophthalma* fly at day time, being especially active just after noon (Brown 1992; Casagrande and Mielke 2000, 2003).

According to Casagrande (2004), *Dasyophthalma* includes four species and five subspecies: the type species, *Dasyophthalma rusina* (Godart, [1824]) — (with three subspecies, *D. r. rusina* (Godart, [1824]), *D. r. delanira* Hewitson, 1862 and *D. r. principesa* Stichel, 1904) — *Dasyophthalma geraensis* Rebel, 1922, *Dasyophthalma creusa* (Hübner, [1821]) — (with two subspecies, *D. c. creusa* (Hübner, [1821]) and *D. c. baronesa* Stichel, 1904) — and *Dasyophthalma vertebralis* Butler, 1869. In previous studies concerning the phylogenetic relationships of *Dasyophthalma* (Penz 2009; Matos-Maravi et al. 2021) two well supported clades were recovered, the first including *D. creusa* + *D. vertebralis* (hereafter “*creusa*-group”) and the second including *D. rusina* + *D. geraensis* (hereafter “*rusina*-group”) (clade names following Penz 2009).

Among the taxa included in the “*rusina*-group”, *D. r. delanira* is quite distinct from all other species and subspecies by both size and coloration of wing marks, showing a distinct pattern of postmedial bands in dorsal forewings. In addition, this subspecies is known from only three localities in the Rio de Janeiro State, restricted to areas above 1200 meters of altitude (Casagrande and Mielke 2008; Penz 2009; Freitas et al. 2018a). Due to these differences, Penz (2009) in her morphological revision of the genus suggested that the status of *D. r. delanira* should be verified.

Very little natural history information, such as morphology of the immature stages and host plants, is available for the genus, and this is basically restricted to the two more common and widespread species, *D. creusa* and *D. rusina* (Casagrande et al. 2000, 2003). This results in a knowledge gap for the three taxa in this group that are included in the Brazilian list of threatened fauna, namely *D. r. delanira*, *D. geraensis*, and *D. vertebralis*, (Casagrande and Mielke 2008; Freitas et al. 2018a, 2018b, 2018c; MMA 2022).

Thus, the present study aims to provide for the first time a phylogenetic assessment of all *Dasyophthalma* species based on molecular data. In addition, the taxonomic status of *D. r. delanira* is revised based on morphological and molecular data. Furthermore, up-to-date distributional data and conservation aspects of the threatened species from the genus are presented and discussed.

2. Methods

2.1. Examined specimens and compiled data

Adult specimens of *D. r. delanira* (Figs 1A, 2), *D. geraensis* (Fig. 1B) and *D. vertebralis* (Fig. 1F, G, Supplementary Material 1) were examined from the following collections: **AMNH**, American Museum of Natural History, New York, New York, USA; **CEIOC**, Coleção Entomológica do Instituto Oswaldo Cruz, Instituto Oswaldo Cruz, Rio de Janeiro, Rio de Janeiro, Brazil; **CGCM**, Carlos Guilherme Costa Mielke research collection, Ponta Grossa, Paraná, Brazil; **DZUP**, Coleção Entomológica Pe. Jesus de Santiago Moure, Universidade Federal do Paraná, Curitiba, Paraná, Brazil; **FLMNH**, McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, University of Florida, Gainesville, Florida, USA; **MCZ**, Museum of Comparative Zoology, Cambridge, Massachusetts, USA; **MNRJ**, Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil; **MZUSP**, Museu de Zoologia, Universidade de São Paulo, São Paulo, Brazil; **NBCN**, Naturalis Biodiversity Center, Leiden, The Netherlands; **NHMUK**, The Natural History Museum, London, UK; **NHMW**, Naturhistorisches Museum Wien, Vienna, Austria; **NHRS**, Swedish Museum of Natural History, Stockholm, Sweden; **RBINS**, Royal Belgian Institute of Natural Sciences, Brussels, Belgium; **USNM**, Smithsonian Institution National Museum of Natural History, Washington, D.C., USA; **ZMUC**, Zoological Museum, University of Copenhagen, Copenhagen, Denmark; **ZUEC**, Coleção Zoológica do Museu de Diversidade Biológica da Universidade Estadual de Campinas, Campinas, São Paulo, Brazil (Table 1).

Data from some collections were compiled from the Global Biodiversity Information Facility (GBIF) (<https://www.gbif.org>). Data from field observations were also compiled and divided in two categories: with and without photographs. Photographs of live specimens were also searched and found on two websites, iNaturalist (www.inaturalist.org), and Facebook (www.facebook.com), which provided useful data on the geographical distribution of these species. Data without photographs came from butterfly specialists (personal communication) (see Table 1). In addition, geographical data were also compiled from Freitas et al. (2018b) and Freitas et al. (2018c) (in these cases, data was not linked to voucher specimens). Geographical data from *D. rusina* (Fig. 1C) and *D. creusa* (Fig. 1D, E, Supplementary Material 1) came exclusively from the datasets of Santos et al. (2018) and Shirai et al. (2019) and also from the iNaturalist website (Fig. 3B, C). The dates of the records of *D. creusa* and *D. rusina* were searched on the iNaturalist website (see Rosa et al. 2023 for details).




Figure 1. Adult *Dasyophthalma* species. **A** *D. delanira* stat. rest.; **B** *D. geraensis*; **C** *D. rusina*; **D**, **E** *D. creusa*, male and female, respectively; **F**, **G** *D. vertebralis*, male and female, respectively; in all pictures, dorsal view at left and ventral view at right. Scale bar = 1 cm.

Table 1. Data for studied individuals of the three threatened species of *Dasyophthalma* from 16 public/private collections (see text or acronyms), literature, and field observations. PA = Protected area (if pertinent).

Taxon	Code	N and Sex	Date	Country or State	Municipality	Site	PA	Altitude (m)
<i>D. delanira</i>	BMNH (type)	1 female	—	—	—	—	—	—
<i>D. delanira</i>	CEIOC	1 male	31.I.1988	RJ	Nova Friburgo	Morro São João, Mury	MCEPA	—
<i>D. delanira</i>	CEIOC	1 male	9.II.1992	RJ	Nova Friburgo	Morro São João, Mury	MCEPA	—
<i>D. delanira</i>	CGCM	1 female	—	RJ	Nova Friburgo	—	—	—
<i>D. delanira</i>	DZUP	3 individuals	30.I.1993	RJ	Nova Friburgo	Morro São João	MCEPA	—
<i>D. delanira</i>	DZUP	1 male	9.II.1992	RJ	Nova Friburgo	Morro São João	MCEPA	—
<i>D. delanira</i>	DZUP	1 individual	14.II.1957	RJ	Nova Friburgo	Mury	MCEPA	—
<i>D. delanira</i>	DZUP	1 individual	13.II.1957	RJ	Nova Friburgo	Mury	MCEPA	—
<i>D. delanira</i>	DZUP	1 female	8.II.1957	RJ	Nova Friburgo	Mury	MCEPA	—
<i>D. delanira</i>	DZUP	10 individuals	9.II.1987	RJ	Nova Friburgo	Pico São João, Mury	MCEPA	1200–1600
<i>D. delanira</i>	DZUP	1 male	—	RJ	Nova Friburgo	—	—	—
<i>D. delanira</i>	MNRJ	3 males	31.I.1988	RJ	Nova Friburgo	Morro São João	MCEPA	—
<i>D. delanira</i>	MNRJ	1 individual	31.I.1988	RJ	Nova Friburgo	P. de São João	MCEPA	1780
<i>D. delanira</i>	ZUEC	1 female	17.III.2023	RJ	Nova Friburgo	RPPN Bacchus	MCEPA	1460
<i>D. delanira</i>	ZUEC	1 female	26.III.2023	RJ	Nova Friburgo	RPPN Bacchus	MCEPA	1460
<i>D. delanira</i>	CGCM	1 individual seen	1996	RJ	Nova Friburgo	Pico do Caledônia	TPSP	—
<i>D. delanira</i>	JS	1 individual seen	10.III.2019	RJ	Nova Friburgo	Pico do Caledônia	TPSP	2050
<i>D. delanira</i>	JMSB	1 individual seen	—	RJ	Cachoeiras de Macacu	GER	BRMEPA	—

Taxon	Code	N and Sex	Date	Country or State	Municipality	Site	PA	Altitude (m)
<i>D. delanira</i>	MP	1 individual seen	9.II.2019	RJ	Nova Friburgo	Pico do Caledônia	TPSP	—
<i>D. geraensis</i>	AMNH	1 male	—	MG	—	—	—	—
<i>D. geraensis</i>	BMNH (type)	1 male	3.II.1916	MG	Virgínia	Fazenda dos Campos	SMEPA	1600
<i>D. geraensis</i>	CEIOC (type)	1 individual	26.I.1921	MG	Virgínia	Fazenda dos Campos	SMEPA	1600
<i>D. geraensis</i>	CEIOC	1 individual	15.I.1920	MG	Virgínia	Fazenda dos Campos	SMEPA	1600
<i>D. geraensis</i>	CEIOC	1 individual	4.II.1916	MG	Virgínia	Fazenda dos Campos	SMEPA	1600
<i>D. geraensis</i>	CEIOC	1 individual	28.I.1923	—	Virgínia	Serra dos Cochos	SMEPA	—
<i>D. geraensis</i>	CEIOC	1 female	20.2.1922	—	Virgínia	Serra dos Cochos	SMEPA	—
<i>D. geraensis</i>	CEIOC	1 individual	15.I.1930	—	—	km 5	—	—
<i>D. geraensis</i>	CEIOC	1 individual	15.I.1936	—	—	km 8	—	—
<i>D. geraensis</i>	CEIOC	1 individual	5.XII.1919	—	—	—	—	—
<i>D. geraensis</i>	CEIOC	1 individual	7.I.1941	—	—	—	—	—
<i>D. geraensis</i>	CGCM	1 male	26.I.1996	MG	Itajubá	—	SMEPA	1400
<i>D. geraensis</i>	DZUP	1 male	17–18.I.2004	MG	Delfim Moreira	15 Km SE	SMEPA	1500–1700
<i>D. geraensis</i>	DZUP	4 individuals	15.II.1984	MG	Delfim Moreira	Barreira de Piquete	SMEPA	1400–1600
<i>D. geraensis</i>	DZUP	1 male	6–11.I.1961	MG	Delfim Moreira	Barreira de Piquete	SMEPA	1500
<i>D. geraensis</i>	DZUP	1 individual	6–11.I.1961	SP	Piquete	Barreira de Piquete	SMEPA	1500
<i>D. geraensis</i>	DZUP	1 male	—	MG	Delfim Moreira	—	SMEPA	—
<i>D. geraensis</i>	DZUP	1 male	—	MG	Virgínia	Fazenda dos Campos	SMEPA	1500
<i>D. geraensis</i>	DZUP (type)	1 male	15.I.1920	MG	Virgínia	Fazenda dos Campos	SMEPA	1600
<i>D. geraensis</i>	DZUP	1 individual	6.II.1919	MG	Virgínia	Fazenda dos Campos	SMEPA	1600
<i>D. geraensis</i>	DZUP	1 individual	18.II.1916	MG	Virgínia	Fazenda dos Campos	SMEPA	1600
<i>D. geraensis</i>	DZUP	1 individual	26.I.1923	MG	Virgínia	Serra dos Cochos	SMEPA	—
<i>D. geraensis</i>	DZUP	1 individual	II.1922	MG	—	Sul de Minas	—	—
<i>D. geraensis</i>	DZUP	1 individual	27.XII.1929	RJ	Itatiaia	Estação biológica	INP	1200
<i>D. geraensis</i>	DZUP	1 individual	12.I.1973	RJ	Itatiaia	INP	INP	1400
<i>D. geraensis</i>	DZUP	1 individual	17.I.1979	RJ	Itatiaia	INP	INP	1750
<i>D. geraensis</i>	DZUP	2 individuals	22.I.1967	RJ	Itatiaia	INP	INP	—
<i>D. geraensis</i>	DZUP	1 individual	12.I.1973	RJ	Itatiaia	INP	INP	1400
<i>D. geraensis</i>	DZUP	1 individual	21.I.1969	RJ	Itatiaia	INP	INP	1300
<i>D. geraensis</i>	DZUP	1 female	6.2.1919	RJ	Itatiaia	—	INP	—
<i>D. geraensis</i>	DZUP	1 individual	—	RJ	Itatiaia	—	INP	—
<i>D. geraensis</i>	DZUP	1 individual	22.III.1969	RJ	—	Serra do Itatiaia	INP	1400
<i>D. geraensis</i>	DZUP	1 individual	15.I.1969	RJ	—	Serra do Itatiaia	INP	1650
<i>D. geraensis</i>	FLMNH	1 male	21.II.1922	ES	Castelo*	—	—	—
<i>D. geraensis</i>	FLMNH	1 male	21.III.1972	RJ	Itatiaia	—	INP	—
<i>D. geraensis</i>	FLMNH	1 male	13–14.I.1973	RJ	—	INP	INP	1000–2000
<i>D. geraensis</i>	MCZ	1 individual	—	MG	—	—	—	—
<i>D. geraensis</i>	MNRJ	2 individuals	—	MG	Passa Quatro	—	SMEPA	—
<i>D. geraensis</i>	MNRJ	1 individual	—	MG	Virgínia	Fazenda dos Campos	SMEPA	—
<i>D. geraensis</i>	MNRJ	1 individual	—	MG	Virgínia	Fazenda dos Campos	SMEPA	1500
<i>D. geraensis</i>	MNRJ	3 males	31.I.1970	RJ	Itatiaia	INP	INP	1650
<i>D. geraensis</i>	MNRJ	1 male	20.I.1953	RJ	Itatiaia	—	INP	—
<i>D. geraensis</i>	MZUSP	1 male	—	MG	Virgínia	Fazenda dos Campos	SMEPA	1500
<i>D. geraensis</i>	MZUSP	1 male	15.I.1936	RJ	Itatiaia	Itatiaia, km 11	INP	—
<i>D. geraensis</i>	MZUSP	1 male	—	RJ	Itatiaia	—	INP	—
<i>D. geraensis</i>	NHFW	1 male (type)	—	MG	Passa Quatro	—	SMEPA	—
<i>D. geraensis</i>	USNM	2 individuals	26.II.	MG	Virgínia	Fazenda dos Campos	SMEPA	—
<i>D. geraensis</i>	USNM	1 individual	—	MG	—	—	—	—
<i>D. geraensis</i>	ZUEC	1 male	30.II.1969	—	—	—	—	—
<i>D. geraensis</i>	ZUEC	2 males	20.I.1969	RJ	Itatiaia	INP, above Maromba	INP	1400–2000
<i>D. geraensis</i>	ZUEC	2 males	3.II.1968	RJ	Itatiaia	INP, above Maromba	INP	1600
<i>D. geraensis</i>	ZUEC	4 males	31.I.1970	RJ	Itatiaia	INP, above Maromba	INP	1400–1700
<i>D. geraensis</i>	ZUEC	1 female	20.I.1969	RJ	Itatiaia	INP, above Maromba	INP	1400–2000
<i>D. geraensis</i>	ZUEC	1 male	3.II.1968	RJ	Itatiaia	INP, above Maromba	INP	1400–1700
<i>D. geraensis</i>	CGCM	1 individual seen	2010	SP	Piquete	Pico dos Marins	SMEPA	—

Taxon	Code	N and Sex	Date	Country or State	Municipality	Site	PA	Altitude (m)
<i>D. geraensis</i>	CRSS	1 individual seen	11.III.2019	MG	Baependi	SPSP	SPSP	1777
<i>D. geraensis</i>	KSB Jr.	1 female seen	15–16. II.1988	MG	Delfim Moreira	11 km SE. of Itajubá	SMEPA	—
<i>D. geraensis</i>	KSB Jr.	1 individual seen	—	MG	Itamonte	—	INP	—
<i>D. geraensis</i>	KSB Jr.	1 individual seen	15.II.1983	RJ	Itatiaia	PNI, Maromba to Macieiras	INP	—
<i>D. geraensis</i>	Freitas et al. 2018b	—	—	SP	Campos do Jordão	CJSP	CJSP	—
<i>D. vertebralis</i>	BMNH (type)	1 male	—	PA*	—	—	—	—
<i>D. vertebralis</i>	CEIOC	2 females	II–III.1948	ES	Sooretama	Parque Sooretama, Cupido	SBR	—
<i>D. vertebralis</i>	DZUP	1 male	—	ES	—	—	—	—
<i>D. vertebralis</i>	DZUP	1 female	—	ES	—	—	—	—
<i>D. vertebralis</i>	DZUP	1 male	—	Peru*	Iquitos	—	—	—
<i>D. vertebralis</i>	DZUP	1 female	—	Peru*	Iquitos	—	—	—
<i>D. vertebralis</i>	MCZ	1 male	—	ES	—	—	—	—
<i>D. vertebralis</i>	MCZ	1 female	—	ES	—	—	—	—
<i>D. vertebralis</i>	MNRJ	1 individual	—	ES	—	—	—	—
<i>D. vertebralis</i>	MNRJ	1 male	—	—	—	—	—	—
<i>D. vertebralis</i>	MNRJ	1 female	—	Peru*	Iquitos	—	—	—
<i>D. vertebralis</i>	MNRJ	1 individual	—	Peru*	Iquitos	—	—	—
<i>D. vertebralis</i>	MZUSP	1 female	—	AM*	—	E. Amazonas	—	—
<i>D. vertebralis</i>	MZUSP	1 female	—	ES	—	—	—	—
<i>D. vertebralis</i>	MZUSP	2 males	—	ES	—	—	—	—
<i>D. vertebralis</i>	NBCN	1 individual	—	—	—	Zuid Amerika	—	—
<i>D. vertebralis</i>	NBCN	1 individual	—	—	—	Zuid Amerika	—	—
<i>D. vertebralis</i>	NHMD	1 male	—	ES	—	—	—	—
<i>D. vertebralis</i>	NHMD	1 male	—	—	—	—	—	—
<i>D. vertebralis</i>	NHMD	1 female	—	—	—	—	—	—
<i>D. vertebralis</i>	NHRS (type)	1 female	—	ES	—	—	—	—
<i>D. vertebralis</i>	RBINS	1 male	—	—	—	—	—	—
<i>D. vertebralis</i>	USNM	1 female	—	ES	Santa Leopoldina	Leopoldina	—	—
<i>D. vertebralis</i>	USNM	1 male	—	—	—	S. Braz.	—	—
<i>D. vertebralis</i>	Freitas et al. 2018c	—	—	ES	Alegre	—	—	—
<i>D. vertebralis</i>	Freitas et al. 2018c	—	—	ES	Muqui	—	—	—
<i>D. vertebralis</i>	Freitas et al. 2018c	—	—	ES	Santa Teresa	—	—	—
<i>D. vertebralis</i>	Freitas et al. 2018c	—	—	MG	Teófilo Otoni	—	—	—

* Individuals possibly mislabeled. Abbreviations for Brazilian states: AM = Amazonas, ES = Espírito Santo, MG = Minas Gerais, PA = Pará, RJ = Rio de Janeiro, SP = São Paulo; Protected areas: BRMEPA = Bacia do Rio Macacu Environmental Protection Area, CJSP = Campos do Jordão State Park, GER = Guapiaçu Ecological Reserve, INP = Itatiaia National Park, MCEPA = Macaé de Cima Environmental Protection Area, SBR = Sooretama Biological Reserve, SMEPA = Serra da Mantiqueira Environmental Protection Area, SPSP = Serra do Papagaio State Park, TPSP = Três Picos State Park; Unpublished data: CGCM = Carlos Guilherme Costa Mielke, CRSS = Carlos Roberto Silva Silva, JS = Jalmirez Silva, JMSB = Jorge Manuel Saraiva Bizarro, MP = Max Peters, KSB Jr. = Keith Spalding Brown Jr.




Figure 2. Adult of *Dasyophthalma delanira* stat. rest. in Nova Friburgo, Rio de Janeiro state, Brazil. **A** Adult with closed wings perched on a leaf; **B** adult with open wings perched on a leaf (photos courtesy of Max Torres Peters).

2.2. Geographical range (area of occupancy and extent of occurrence)

Geographical range (extent of occurrence EOO and area of occupancy AOO) was estimated based on all known sites for each species of *Dasyophthalma*. The EOO is the area contained within the shortest continuous imaginary boundary that includes all known distribution points of a species, and the AOO is the area within its EOO that is really occupied by a taxon (IUCN 2012; IUCN Standards and Petitions Committee 2022). Both EOO and AOO were estimated using the online open-source program GeoCAT (Geospatial Conservation Assessment Tool, available at <http://geocat.kew.org>) (Bachman et al. 2011). As recommended by IUCN for AOO analyses, a 2 km grid (cell area of 4 km²) was used (IUCN 2012; IUCN Standards and Petitions Committee 2022). Doubtful geographical data were not used in these analyses.

2.3. DNA purification, amplification, and sequencing

The mitochondrial cytochrome c oxidase subunit I (COI) and the nuclear glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and ribosomal protein S5 (RpS5) genes were selected for this study.

Total genomic DNA was purified from one or two legs per individual using the DNeasy Blood & Tissue Kit protocol (QIAGEN, Düsseldorf, Germany). DNA was stored in TE buffer at −20°. PCR amplification of cytochrome c oxidase subunit I (COI) partial sequences (ca. 658 bp) were performed using the LCO 1490 (Forward; 5' GGTC AACAAATCATAAAGATATTGG) and HCO 2198 (Reverse; 5' TAAACTTCAGGGTGACCAAAAAATCA) primers by Folmer et al. (1994). Reactions were done in 26 µL µL and contained 15,6 µL H₂O Milli-Q, 2,5 µL buffer solution (500 Mm KCl), 0,4 µL of dNTP (25 Mm), 2,0 µL of MgCl₂ (2 Mm), 2,5 µL of DMSO 5%, 0,5 µL of each initiator (10 Mm), 0,2 µL of Taq DNA-Polymerase

(Promega, Madison, WI, USA) (1 U), and 2,0 µL of DNA extracted. The amplification program included an initial denaturation step at 94°C for 2 min, followed by 34 cycles of denaturation at 94°C for 45s, annealing at 45°C for 45s, and polymerization at 72°C for 1 min, followed by an extension step at 72°C for 5 min (Silva-Brandão et al. 2005). PCR products were purified of primers and deoxynucleotides with ExoSAP-IT (GE Healthcare, Bucks, UK), and then sequenced by BigDye Kit protocol in an ABI 3500 automated sequencer, with primers used for amplification.

In the case of two specimens of *D. vertebralis* (voucher code: EB-19009 and EB-19033), DNA was extracted from old museum specimens deposited at the Natural History Museums at Stockholm and Copenhagen, respectively. The DNA extraction was carried out using QIAamp DNA MicroKit (QIAGEN®, USA) protocol, adapted with columns from MinElute PCR Purification Kit (QIAGEN®, Düsseldorf, Germany).

The standard steps for NGS library preparation (e.g., end repair, adaptor ligation and fill-in, and indexing PCR) were used, using Blunt-End Illumina Libraries, which consists in single index blunt-end Illumina library construction for ancient and historical samples, following a modified protocol of Meyer and Kircher (2010) as described in Twort et al. (2021). The pooled libraries were sent to sequencing at National Genomics Infrastructure (NGI) in Stockholm, Sweden on an Illumina NovaSeq machine.

All libraries were successfully sequenced using Whole Genome Shotgun Sequencing (WGSS) and the raw data were analyzed and each genome has been cleaned and assembled using de novo assembly techniques, as described in Twort et al. (2021). The COI and GAPDH genes were then successfully pooled from the two genomes and incorporated in the matrix.

Sequences generated in this work and from Matos-Maraví et al. (2021) were aligned with the sequences from COI and the nuclear genes GAPDH and RpS5 (Table 2 for accession numbers) using the program Geneious v. 11.1.5 (Kearse et al. 2012).




Figure 3. Distribution map showing the known geographic distribution of *Dasyophthalma* species in Brazil. **A** Distribution of *D. delanira*, *D. geraensis* and *D. vertebralis*; **B** distribution of *D. rusina*; **C** distribution of *D. creusa*.

Table 2. Samples of Satyrinae (Morphini and Brassolini) used in the present study with codes, localities and GenBank/Boldsystems accession numbers for COI, GAPDH and RpS5 sequences.

Taxon	DNA voucher	Locality	COI	GAPDH	RpS5
<i>Morpho helenor</i>	NW127-12	Águas da Prata, São Paulo, Brazil	JN696174	JN696220	JN696282
<i>Bia actorion</i>	PM14-28	Phil De Vries Collection	MK551389	MK551492	MK551469
<i>Bia actorion</i>	NW17842	Xingu, Pará, Brazil	MK551366	—	MK551450
<i>Blepopenis bassus</i>	NW155-6	Floresta Nacional, São Francisco de Paula, Rio Grande do Sul, Brazil	JF508389	MK551479	JF508405
<i>Blepopenis bassus</i>	BC-CGCM-8.908	São Bento do Sapucaí, Brazil	JX215903	—	—
<i>Blepopenis bassus</i>	BC-CGCM-9.339	São Bento do Sapucaí, Brazil	JX215904	—	—
<i>Blepopenis batea</i>	BC-CGCM-18.416	Boca do Mato, Cachoeiras de Macacu, Rio de Janeiro, Brazil	JX215905	—	—
<i>Blepopenis batea</i>	NW155-3	Paranapiacaba, Santo André, São Paulo, Brazil	JF508388	KM013289	JF508404
<i>Blepopenis batea</i>	BPU172	Parque Nacional Serra da Bocaina, São José do Barreiro, São Paulo, Brazil	OQ720958	—	—
<i>Blepopenis catharinae</i>	PM02-09	Lagoa do Peri, Florianópolis, Santa Catarina, Brazil	JF508390	—	JF508406
<i>Orobassolis ornamentalis</i>	PM25-05	Bairro Novo Capivari, Campos do Jordão, São Paulo, Brazil	MK551395	—	MK551473
<i>Orobassolis ornamentalis</i>	BLU-785	Campos do Jordão, São Paulo, Brazil	MK551348	—	—
<i>Orobassolis ornamentalis</i>	BPU-169	Campos do Jordão, São Paulo, Brazil	OQ720956	—	—
<i>Orobassolis ornamentalis</i>	BPU-202	Campos do Jordão, São Paulo, Brazil	OQ720963	—	—
<i>Opsiphanes bogotanus</i>	NW118-6	Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JF508393	MK551476	JF508408
<i>Opsiphanes bogotanus</i>	08-SRNP-40116	Sector Rincon Rain Forest, Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JQ537131	—	—
<i>Opsiphanes cassiae</i>	PM06-11	Misahualli, Napo, Ecuador	KM012972	—	KM013196
<i>Opsiphanes cassina</i>	NW118-5	Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JF508392	—	—
<i>Opsiphanes cassina</i>	YB-BC124435	Barro Colorado, Panamá, Panamá	KP849172	—	—
<i>Opsiphanes invirae</i>	MACN-Bar-Lep-et-02127	Parque Nacional Iguazú, Departamento de Iguazu, Misiones, Argentina	MF547146	—	—
<i>Opsiphanes invirae</i>	MACN-Bar-Lep-et-02129	Parque Nacional Iguazú, Departamento de Iguazu, Misiones, Argentina	MF547363	—	—
<i>Opsiphanes quiteria</i>	NW109-10	Area de Conservacion Guanacaste, Guanacaste, Costa Rica	DQ018957	KM013296	EU528451
<i>Opsiphanes quiteria</i>	BLU263	Serra do Japi, Jundiá, São Paulo, Brazil	KX858943	—	—
<i>Opsiphanes quiteria</i>	BLU322	Serra do Japi, Jundiá, São Paulo, Brazil	KX858944	—	—
<i>Opsiphanes tamarindi</i>	08-SRNP-56847	Sector Mundo Nuevo, Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JQ538545	—	—
<i>Opsiphanes tamarindi</i>	07-SRNP-33627	Sector Pitilla, Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JQ537147	—	—
<i>Opsiphanes tamarindi</i>	NW118-3	Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JF508391	—	JF508407
<i>Dynastor datus</i>	06-SRNP-60108	Sector Mundo Nuevo, Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JQ548386	—	—
<i>Penetes pamphanis</i>	NW155-5	Trilha Pedra de Amolar, Maquiné, Rio Grande do Sul, Brazil	KM012973	KM013304	KM013219
<i>Dasyophthalma creusa</i>	NW126-4	Serra do Japi, Jundiá, São Paulo, Brazil	EU528318	EU528387	EU528431
<i>Dasyophthalma delanira</i>	BC-CGCM-7.561	Pico São João, Nova Friburgo, Rio de Janeiro, Brazil	JX215914	—	—
<i>Dasyophthalma delanira</i>	BC-DZ-9.911	Pico São João, Mury, Nova Friburgo, Rio de Janeiro, Brazil	CGCM377-08	—	—
<i>Dasyophthalma geraensis</i>	BC-CGCM-3.035	Itajuba, Minas Gerais, Brazil	JX215915	—	—
<i>Dasyophthalma geraensis</i>	BC-DZ-9.903	Delfim Moreira, Minas Gerais, Brazil	CGCM381-08	—	—
<i>Dasyophthalma rusina</i>	BC-CGCM-9.492	Boca do Mato, Cachoeiras de Macacu, Rio de Janeiro, Brazil	JX215917	—	—
<i>Dasyophthalma rusina</i>	BC-CGCM-20.515	Conceição dos Ouros, Minas Gerais, Brazil	JX215916	—	—

Taxon	DNA voucher	Locality	COI	GAPDH	RpS5
<i>Dasyophthalma rusina</i>	BC-DZ-9.887	Pico São João, Mury, Nova Friburgo, Rio de Janeiro, Brazil	CGCM380-08	—	—
<i>Dasyophthalma rusina</i>	BC-DZ-9.919	Rio Natal, São Bento do Sul, Santa Catarina, Brazil	CGCM382-08	—	—
<i>Dasyophthalma rusina</i>	NW155-8	Paranapiacaba, Santo André, São Paulo, Brazil	MK551359	MK551480	MK551443
<i>Dasyophthalma rusina</i>	BPU187	Vossoroca, Tijucas do Sul, Paraná, Brazil	OQ720959	—	—
<i>Dasyophthalma rusina</i>	BPU191	Parque Natural Municipal do Trabalhú, Pindamonhangaba, São Paulo, Brazil	OQ720960	—	—
<i>Dasyophthalma rusina</i>	BPU192	Serra do Japi, Jundiaí, São Paulo, Brazil	OQ720961	—	—
<i>Dasyophthalma rusina</i>	BPU193	Parque Nacional do Caparaó, Alto Caparaó, Minas Gerais, Brazil	OQ720962	—	—
<i>Dasyophthalma rusina</i>	BPU208	Parque Nacional do Itatiaia, Itatiaia, Rio de Janeiro, Brazil	OQ720964	—	—
<i>Dasyophthalma rusina</i>	BPU268	RPPN Bacehus, Nova Friburgo, Rio de Janeiro, Brazil	OQ720965	—	—
<i>Dasyophthalma vertebralis</i>	EB19-009	Espírito Santo, Brazil	OQ720966	OQ722368	—
<i>Dasyophthalma vertebralis</i>	EB19-033	Espírito Santo, Brazil	OQ720967	OQ722369	—
<i>Catoblepia amphirhoe</i>	MACN-Bar-Lep-ct-01007	Parque Nacional Iguazú, Departamento de Iguazú, Misiones, Argentina	MF546750	—	—
<i>Catoblepia amphirhoe</i>	MACN-Bar-Lep-ct-01178	Parque Nacional Iguazú, Departamento de Iguazú, Misiones, Argentina	MF546831	—	—
<i>Catoblepia berecynthia</i>	NW155-14	Trap 25 understory, Garza Cocha, Sucumbios, Ecuador	MK551355	—	MK551439
<i>Catoblepia berecynthia</i>	NW178-38	Xingu, Pará, Brazil	MK551363	MK551483	MK551447
<i>Catoblepia berecynthia</i>	YB-BC164698	Gamboia, Colón, Panamá	KP848752	—	—
<i>Catoblepia berecynthia</i>	YB-BC164818	Gamboia, Colón, Panamá	KP848753	—	—
<i>Catoblepia xanthus</i>	BLU024	Itacoatiara, Amazonas, Brazil	KX858937	—	—
<i>Catoblepia xanthus</i>	BLU025	Itacoatiara, Amazonas, Brazil	KX858938	—	—
<i>Catoblepia xanthus</i>	NW155-13	Trap 14 understory, Garza Cocha, Sucumbios, Ecuador	JF508395	—	JF508410
<i>Catoblepia soranus</i>	NW178-39	Xingu, Pará, Brazil	MK551364	—	MK551448
<i>Catoblepia verstitincta</i>	NW178-40	Xingu, Pará, Brazil	MK551365	MK551484	MK551449
<i>Selenophanes cassiope</i>	YPH-0585	Foz do Iguaçu, Paraná, Brazil	KX858949	—	—
<i>Selenophanes cassiope</i>	YPH-0587	Foz do Iguaçu, Paraná, Brazil	KX858948	—	—
<i>Selenophanes orgetorix</i>	NW109-15	Area de Conservacion Guanacaste, Guanacaste, Costa Rica	DQ338754	—	EU528427
<i>Ooptera aorsa</i>	MACN-Bar-Lep-ct-01206	Parque Nacional Iguazú, Departamento de Iguazú, Misiones, Argentina	MF545789	—	—
<i>Ooptera aorsa</i>	MACN-Bar-Lep-ct-01208	Parque Nacional Iguazú, Departamento de Iguazú, Misiones, Argentina	MF545409	—	—
<i>Ooptera aorsa</i>	NW137-21	Cotia, São Paulo, Brazil	KM012970	—	MK551437
<i>Ooptera syme</i>	BLU511	Serra do Japi, Jundiaí, São Paulo, Brazil	KX858935	—	—
<i>Ooptera syme</i>	NW126-3	Serra do Japi, Jundiaí, São Paulo, Brazil	EU528323	EU528403	EU528450
<i>Ooptera arsippe</i>	NN33	Peru	MK551352	—	MK551436
<i>Narope cyllabarus</i>	PM01-24	Três Lagoas, Mato Grosso do Sul, Brazil	KM012968	KM013295	KM013211
<i>Narope cyllene</i>	NW127-27	Extrema, Minas Gerais, Brazil	DQ338755	EU528401	EU528447
<i>Caligo brasiliensis</i>	10-SRNP-21447	Sector Del Oro, Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JQ539210	—	—
<i>Caligo brasiliensis</i>	10-SRNP-21446	Sector Del Oro, Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JQ539211	—	—
<i>Caligo illioneus</i>	06-SRNP-2481	Area de Conservacion Guanacaste, Alajuela, Costa Rica	JQ548328	—	—
<i>Caligo illioneus</i>	MACN-Bar-Lep-ct-02571	Parque Nacional Rio Pilcomayo Seccional Estero Poi, Pilcomayo, Formosa, Argentina	MF547002	—	—

Taxon	DNA voucher	Locality	COI	GAPDH	RpS5
<i>Caligo illioneus</i>	PM06-17	Puente Chinchavito, Leoncio Prado, Peru	MK551377	MK551488	MK551461
<i>Caligo uranus</i>	PM06-15	Quintana Roo, Othon P. Blanco, Mpio, Bacalar, Mexico	MK551375	MK551487	MK551459
<i>Eryphanis aescacus</i>	MAL-02690	La Union, Rio Hondo, Quintana Roo, Mexico	GU659596	—	—
<i>Eryphanis aescacus</i>	MAL-02692	Chetumal, Quintana Roo, Mexico	GU659598	—	—
<i>Eryphanis reevesii</i>	MACN-Bar-Lep-ct-00287	Parque Nacional Iguazú, Departamento de Iguazu, Misiones, Argentina	MF546004	—	—
<i>Eryphanis lycomedon</i>	INB0004269656	Nogal, Puerto Viejo, Sarapiquí, Heredia, Costa Rica	ASARD2471-12	—	—
<i>Eryphanis lycomedon</i>	16-SRNP-31641	Area de Conservacion Guanacaste, Alajuela, Costa Rica	BLPAA5850-17	—	—
<i>Eryphanis lycomedon</i>	PM06-06	Pichincha, Tinalandia, Peru	MK551368	MK551485	MK551452
<i>Eryphanis lycomedon</i>	PM06-07	Pichincha, Tinalandia, Peru	MK551369	—	MK551453
<i>Eryphanis automedon</i>	NW155-18	CICRA, Peru	KM012950	KM013287	KM013193
<i>Brassolis sophorae</i>	NW122-21	Campinas, São Paulo, Brazil	EU528314	GQ357384	EU528425
<i>Brassolis sophorae</i>	BLU617	UNICAMP, Campinas, São Paulo, Brazil	OQ720957	—	—
<i>Brassolis sophorae</i>	BC-DZ-9.863	Cambé, Paraná, Brazil	CGCM373-08	—	—
<i>Brassolis isidrochaconi</i>	MHMYJ1180-11	Sector Rincon Rain Forest, Area de Conservacion Guanacaste, Alajuela, Costa Rica	JQ539205	—	—
<i>Brassolis isidrochaconi</i>	07-SRNP-6120	Sector Santa Maria, Area de Conservacion Guanacaste, Guanacaste, Costa Rica	JQ536912	—	—

Table 3. Best-fit substitution models by partition derived from ModelFinder on IQTree 2.1.2.

Partition	Name	Best fit-model
1	COI_pos1	TIM2+F+G4
2	COI_pos2	TIM2+F+I+G4
3	COI_pos3 GAPDH_pos3 RPS5_pos3	TPM2+F+I+G4
4	GAPDH_pos1 RPS5_pos1	TPM2+F+G4
5	GAPDH_pos2_RPS5_pos2	TN+F+I+G4

2.4. Phylogenetic analysis

The concatenated matrix comprised three genes (mitochondrial COI and nuclear GAPDH and RpS5) and 88 specimens of 40 Satyrinae species (39 *Brassolini* and one *Morphini*, used to root the tree), representing all five *Dasyophthalma* species, with a total of 2,776 base pairs (COI – 1475bp; GAPDH – 691bp; RpS5 – 610bp) (see Table 2 for voucher numbers and information on the newly generated sequences). Data was partitioned by codon position and model selection (see Table 3 for details) was done using ModelFinder (Kalyaanamoorthy et al. 2017) with edge-linked partition model + FreeRate heterogeneity in IQ-TREE v. 2.1.2. (Minh et al. 2020) on CIPRES portal (Miller et al. 2010). The concatenated data set with aligned sequences is provided in the Supplementary Material (Supplementary Material 2).

The maximum likelihood tree was inferred using 10 independent likelihood searches in IQ-TREE v. 2.1.2, and support was calculated using the ultrafast bootstrap (UFBoot) (Hoang et al. 2018), with 2000 replications. In addition to assess node support through 1000 replications of Shimodaira Hasegawa-like approximate Likelihood Ratio Test (SH-aLRT) (-aLRT 1000) (Guindon et al. 2010; Hoang et al. 2018) and Approximate Bayes test (aBayes) (Anisimova et al. 2011). The tree associated with the highest likelihood score (LnL = -16552.4484) was rooted as described above.

The genetic distances (Table 4) among species of *Dasyophthalma* were determined by using the nucleotide substitution model Kimura-2-parameters (Kimura 1980) by using the program MEGA X v.10.1.7 (Kumar et al. 2018).

3. Results

3.1. Compiled data

A total of 28 adult specimens of *D. r. delanira*, 67 of *D. geraensis* and 26 of *D. vertebralis* were found in 16 public/private collections (Table 1). Data from field observations “without photographs” (pers. comm.) are two of *D. r. delanira*, and four of *D. geraensis*. Data “with photographs” came from Facebook, two of *D. r. delanira*, one of *D. geraensis*. Literature data (Freitas et al. 2018b,c) are three of *D. vertebralis* and 1 of *D. geraensis*. Data of *D.*




Figure 4. Phylogenetic relationships of *Dasyophthalma* species and other Satyrinae taxa based on COI, GAPDH and RpS5 genes and obtained by a maximum likelihood analysis. Numbers near the nodes are SH-aLRT/aLRT/UFBoot support values.

creusa are 41 from datasets and 76 from iNaturalist and for *D. rusina* 34 from datasets and 27 from iNaturalist website.

3.2. Phylogenetic assessment and taxonomy

The obtained molecular phylogeny (Fig. 4) recovered the genus *Bia* (subtribe Biina) as the sister group of all remaining Brassolini (subtribe Brassolina). In the subtribe Brassolina, five main clades were identified: 1) *Narope* + *Ooptera* (hereafter “*Ooptera* clade”) as the sister group of all remaining Brassolina; 2) *Dynastor* + *Dasyophthalma* (“*Dasyophthalma* clade”); 3) *Caligo* + *Eryphanis* (“*Caligo* clade”); 4) the genus *Brassolis* (“*Brassolis* clade”), as the sister group of 5) a large clade with all remaining genera (“*Opsiphanes* clade”). All brassoline genera were recovered as monophyletic, including *Dasyoph-*

thalma, which have *Dynastor darius* as sister group. The two species groups of *Dasyophthalma* previously defined (Penz 2009) were also recovered as sister to each other: (1) the “*creusa* group”, including *D. creusa* + *D. vertebralis*, and (2) the “*rusina*-group”, including *D. delanira stat. rest.* + *D. geraensis*, this clade sister to *D. rusina* (Fig. 4).

Based on the COI, the mean K2P distance within species of *Dasyophthalma* was 0.017 (range 0 to 0.03), and the mean among species distance was 0.071 (range 0.02 to 0.10). The average distance between *D. delanira stat. rest.* and *D. rusina* varied from 0.05 to 0.07 (mean 0.056), and between *D. delanira stat. rest.* and *D. geraensis* was 0.02, both above the mean distance within species (Table 4).

In short, the present results recovered *D. delanira stat. rest.* as the sister species of *D. geraensis*, and this clade sister to *D. rusina*. This, combined with the genetic distances validate the decision of reinstating the specific status of the former, and not as a subspecies of *D. rusina*.

Table 4. Pairwise Genetic distances (Kimura 2-parameter distance) among sampled *Dasyophthalma* species.

		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
1	<i>Dasyophthalma vertebralis</i> EB19009																	
2	<i>Dasyophthalma vertebralis</i> EB19033	0.00																
3	<i>Dasyophthalma creusa</i> NW126-4	0.06	0.06															
4	<i>Dasyophthalma delanira</i> BC-CGCM-7.561	0.09	0.09	0.09														
5	<i>Dasyophthalma delanira</i> BC-DZ-9.911	0.09	0.09	0.09	0.00													
6	<i>Dasyophthalma geraensis</i> BC-CGCM-3.035	0.09	0.09	0.10	0.02	0.02												
7	<i>Dasyophthalma geraensis</i> BCDZ9.903	0.09	0.09	0.10	0.02	0.02	0.00											
8	<i>Dasyophthalma rusina</i> BC-CGCM-9.492	0.09	0.09	0.08	0.05	0.05	0.05	0.05										
9	<i>Dasyophthalma rusina</i> BC-CGCM-20.515	0.09	0.09	0.09	0.06	0.06	0.06	0.06	0.02									
10	<i>Dasyophthalma rusina</i> BC-DZ-9.887	0.09	0.09	0.08	0.05	0.05	0.05	0.05	0.00	0.02								
11	<i>Dasyophthalma rusina</i> BC-DZ-9.919	0.10	0.10	0.09	0.06	0.06	0.06	0.06	0.03	0.02	0.03							
12	<i>Dasyophthalma rusina</i> NW155_8	0.09	0.09	0.08	0.05	0.05	0.05	0.05	0.02	0.01	0.02	0.01						
13	<i>Dasyophthalma rusina</i> BPU191	0.09	0.09	0.09	0.06	0.06	0.06	0.06	0.02	0.00	0.02	0.02	0.01					
14	<i>Dasyophthalma rusina</i> BPU192	0.09	0.09	0.09	0.06	0.06	0.06	0.06	0.02	0.00	0.02	0.02	0.01	0.00				
15	<i>Dasyophthalma rusina</i> BPU208	0.10	0.10	0.09	0.07	0.07	0.06	0.06	0.02	0.00	0.02	0.02	0.02	0.00	0.00			
16	<i>Dasyophthalma rusina</i> BPU268	0.09	0.09	0.08	0.05	0.05	0.05	0.05	0.00	0.02	0.00	0.03	0.02	0.02	0.02	0.02		
17	<i>Dasyophthalma rusina</i> BPU187	0.10	0.10	0.09	0.06	0.06	0.06	0.06	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	
18	<i>Dasyophthalma rusina</i> BPU193	0.10	0.10	0.09	0.05	0.05	0.06	0.06	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.04	0.03	0.03

3.3. Distribution and habitat

Based on the available geographical data, the genus *Dasyophthalma* is endemic from Brazil and has its distribution mostly in the Atlantic Forest domain, with only a few records in the Cerrado domain (Fig. 3). The most restricted species is *D. delanira* **stat. rest.**, known from only four localities of well-preserved montane forest at altitudes above 1200 meters of two municipalities (Nova Friburgo and Cachoeiras de Macacu) in the Rio de Janeiro State (Fig. 3A, Table 1). Another high-altitude montane forest species, *D. geraensis*, was reported in 11 localities above 1000 meters of altitude along the Mantiqueira mountain range, in the states of Minas Gerais, São Paulo, and Rio de Janeiro (Figs 3A, 5C, D; Table 1). There is a single record of *D. geraensis* for the municipality of Castelo in the State of Espírito Santo that needs to be confirmed and could represent a mistake. The most enigmatic of the species of the genus is *D. vertebralis*; although the available data suggests a relatively wide distribution, in lowland Tableland forests in six localities in the states of Minas Gerais and Espírito Santo (Fig. 3A, Supplementary Material 1, Table 1), no individuals were observed since 1948, when two individuals were captured in Sooretama Biological Reserve, Espírito Santo State (Supplementary Material 1). In addition, the occurrence of this species in the Brazilian state of Pará, in the Amazon region (mentioned in the original description) and in Iquitos (Peru), are both considered as mislabeled specimens (Freitas et al. 2018c). Two species, *D. creusa* and *D. rusina*, are common and show a wider geographical range, occurring in sympatry in great part of its distribution (Fig. 3B, C). However, *D. rusina* is more widespread and common in altitudes above 500 m, also occurring in forest areas within the Cerrado in Central Brazil (in Minas Gerais state and the Federal District), (Figs 3B, 5A, B), while *D. creusa* is more common on

lowland coastal forests (on all states of Southeastern and South Brazil) (Figs 3C, 5A, B).

3.4. Natural history

Species of *Dasyophthalma* fly during the day, including the warmer hours (from 11:00 to 14:00), a pattern distinct of most Brassolini, which present crepuscular behavior (Brown 1992; Freitas et al. 1997; Casagrande and Mielke 2000, 2003). Adults are commonly observed in an erratic and fast flight in the forest understory, occasionally leaving to forest edges along rivers, dirty roads and trails. Based on all available data and field observations, all species of *Dasyophthalma* are univoltine, with adults flying during the summer months (records are mostly from December to March, with few records in November, April and May) (Table 1; see also Brown 1992; Penz 2009: Appendix 1; data from iNaturalist and personal observations of the authors also indicate the same flight period). Territorial behavior was not observed, and courtship behavior is unknown.

Some species occur in sympatry in several localities, such as *D. creusa* with *D. rusina* (Fig. 5A, B) and *D. creusa* with *D. vertebralis* (Supplementary Material 1). Others occurs in a parapatry as a result of different altitudinal preferences; the most notable example is *D. rusina* and *D. delanira* **stat. rest.**, where the latter occurs in altitudes higher than the former, resulting in a very narrow line of sympatry around 1200–1300 m.

As most Brassolini, all species of *Dasyophthalma* belong to the fruit-feeding guild, and the most common species (i.e., *D. creusa* and *D. rusina*) are usually sampled in studies using traps baited with fermented fruits (e.g. Uehara-Prado et al. 2007), however, the other species in the genus are also potentially attracted to bait traps (Freitas and Marini-Filho, 2011; Freitas et al. 2018a, 2018b,




Figure 5. Habitat of *D. rusina*, *D. creusa* and *D. geraensis* in Itatiaia National Park (INP). **A** Closer view of *D. rusina* and *D. creusa* habitat; **B** general view of *D. rusina* and *D. creusa* habitat; **C** closer view of *D. geraensis* habitat; **D** general view of *D. geraensis* habitat.

2018c). It is worth mentioning that fieldwork to sample *D. vertebralis* using attractive traps was carried out at Sooretama Biological Reserve on March 2020 without success (Supplementary Material 1). Also, a previous three-month sampling of fruit-feeding butterflies in the same area (April, July and August) captured only *D. creusa* in baited traps (Nogueira 2012).

Immature stages and host plants have been described (partially) only for the two most common species, *D. creusa* and *D. rusina*. For *D. geraensis*, its host plant “uricana/uricangas” (*Bactris tormentosa*, Arecaceae) is mentioned in some old studies, but immature stages are unknown (d’Araújo e Silva et al. 1968; Zikán and Zikán 1940; Beccaloni et al. 2008). For *D. creusa*, the fourth and fifth instars and the pupa were described (Casagrande and Mielke 2003) and larvae were fed with *Geonoma schottiana* (Arecaceae) as host plant. Other host plants mentioned in the literature include “taquara” (Poaceae) (Hoffmann 1936), *Astrocaryum aculeatum/vulgare?* (Arecaceae) (Schmith and Hoffmann 1931) and *Bactris* sp. (Arecaceae) (Brown 1992). For *D. rusina*, the fifth instar and pupa were described, and its host plant is *Geonoma schottiana* (Arecaceae) (Casagrande & Mielke 2000). Other reported host plants include *Euterpe edulis* (Arecaceae) (Zikán and Zikán 1940; d’Araújo e Silva et

al. 1968) and *Bambusa* sp. (Poaceae) (d’Araújo e Silva et al. 1968; Brown 1992).

Population data is limited and once again restricted to the two common and widespread species, *D. creusa* and *D. rusina*. Available data using baited traps reveal a large variation in number of sampled individuals in different areas. For example, for *D. creusa*, a minimum of one individual and a maximum of 81 individuals are reported in several short-term studies (Supplementary Material 1). For *D. rusina*, the few short-term studies sampled from one to 11 individuals (Table S1). In the only long-term study, in a montane area in southeastern Brazil, numbers were low, with only 35 individuals of *D. creusa* (30 males and five females) and three individuals of *D. rusina* (one male and two females) captured over 10 years of sampling (Supplementary Material 1).

3.5. Conservation

As previously mentioned, three out of the five species of *Dasyophthalma* are of conservation importance, namely *D. delanira* **stat. rest.**, *D. geraensis* and *D. vertebralis* (MMA 2003, 2014, 2022; Freitas et al. 2018a, 2018b, 2018c). In these three cases, the inclusions on the Bra-

Table 5. Conservation status of all three threatened species of *Dasyophthalma* species along different assessments from 1989 to 2022 (from the Official Brazilian Red List).

Species	1989 ^a	2003 ^b	2014 ^c	2022 ^d	Future ^e
<i>D. delanira</i>	Threatened	Threatened	CR	EN	EN
<i>D. geraensis</i>	Threatened	Threatened	CR	CR	VU
<i>D. vertebralis</i>	Threatened	Threatened	CR-PEX	CR-PEX	DD
<i>D. creusa</i>	—	—	—	LC*	LC
<i>D. rusina</i>	—	—	—	LC*	LC

CR = Critically endangered, CR-PEX = Critically endangered, probably extinct; EN = Endangered; DD = Data Deficient; LC = Least Concern; LC* = Least Concern (Freitas et al. 2022a,b; Freitas AVL, Brant A, Rosa AHB, Ribeiro DB, Barbosa EP, Dias FMS, Bizarro JMS, Prado UM, Cardoso MZ, Filho OJM, Taumaturgo TZB (2022a). *Dasyophthalma rusina* (Godart, [1824]), Apr/2018. Sistema de Avaliação do Risco de Extinção da Biodiversidade – SALVE. <https://salve.icmbio.gov.br/salve/> and Freitas AVL, Brant A, Rosa AHB, Ribeiro DB, Barbosa EP, Dias FMS, Bizarro JMS, Prado UM, Cardoso MZ, Filho OJM, Taumaturgo TZB (2022b) *Dasyophthalma creusa* (Hübner, [1821]), Apr/2018. Sistema de Avaliação do Risco de Extinção da Biodiversidade – SALVE. <https://salve.icmbio.gov.br/salve/>). ^a = IBAMA 1989; ^b = MMA 2003; ^c = MMA 2014; ^d = MMA 2022; ^e = Unpubl. Data.

Table 6. Geographic range of *Dasyophthalma* species

Species	EOO (km ²)	AOO (km ²)
<i>D. delanira</i>	89.90	16
<i>D. geraensis</i>	2,336.36	40
<i>D. vertebralis</i>	30,556.44	24
<i>D. creusa</i>	662,297.69	408
<i>D. rusina</i>	917,612.14	224

EOO = Extent of occurrence; AOO = Area of occupancy.

zilian Red list were all based on the IUCN B criterion (geographical range), since there is no data on population trends for them. The changes in the conservation status of each three species over time are shown in Table 5.

In addition of being on the national Red List, these three species were/are also listed on several regional Red lists, such as for the state of Minas Gerais (COPAN 1996; Casagrande et al. 1998; COPAN 2010), Espírito Santo (Espírito Santo 2005; Azevedo et al. 2007; Fraga et al. 2019), Rio de Janeiro (SEMA 1998; Otero et al. 2000), and São Paulo (São Paulo 2008, 2010, 2014, 2018).

However, for at least two threatened species of *Dasyophthalma* the situation is optimistic, since several known populations are located inside protected areas (PAs) (IUCN Protected Area Management Categories I–VI (I–V = fully protected areas, and VI “Protected area with sustainable use of natural resources”, see Dudley 2008)). For *D. delanira* **stat. rest.**, all four known localities of occurrence are inside three protected areas (two in two PAs VI category and one in a fully PA) and for *D. geraensis*, all 11 localities of occurrence are inside four protected areas (seven in one PA VI category and four in three fully PAs). For *D. vertebralis*, however, only one out of the six known localities of occurrence is inside a fully protected area (Table 1).

Based on geographic distribution records, the geographical ranges (EOO and AOO) were estimated and are presented in Table 6. Based only on the thresholds in km² of the IUCN “B” criterion (geographic range), *D. delanira* **stat. rest.** and *D. geraensis* present values of EOO that would keep them as threatened by the subcriterion “B1”,

the others present values greater than 30,000,00 km² and are not able to figure under “B1” (Table 6). However, the AOO threshold values would include all species under the sub-criterion “B2” (Table 6).

4. Discussion

4.1. Phylogeny and taxonomy

The present phylogeny recovered five main clades that exactly correspond to the ones obtained by Matos-Maraví et al. (2021) based also on molecular data. Accordingly, the “*Dasyophthalma* clade” corresponds to ‘clade A’ of Matos-Maraví et al. (2021), the “*Opoptera* clade” corresponds to ‘clade B’, the “*Caligo* clade” corresponds to ‘clade C’, the “*Opsiphanes* clade” corresponds to ‘clade D’ and the “*Brassolis* clade” corresponds to an unnamed clade including only the genus *Brassolis*. The main difference is the position of the “*Opoptera* clade”; in the present study this clade is the sister group of all remaining *Brassolina*, a position occupied by the “*Dasyophthalma* clade” (as ‘clade A’) in Matos-Maraví et al. (2021). The results are also somewhat congruent with those presented by Penz et al. (2013), based on morphology. In that study, three out of the five main clades have correspondence with the present study, namely, the “*Caligo* clade”, the “*Brassolis* clade” and the “*Opsiphanes* clade”; the “*Dasyophthalma* clade” was not recovered, with *Dynastor* appearing as sister to *Brassolis* (Penz et al. 2013) and *Dasyophthalma* as sister to *Opoptera* in some analyses. Concerning the genus *Dasyophthalma*, the present results are virtually identical to those obtained in previous morphological and molecular studies, with two well supported clades (Penz 2009; Matos-Maraví et al. 2021), the “*creusa*-group” and the “*rusina*-group” (*sensu* Penz 2009). However, our study is the first to include samples of *D. delanira* **stat. rest.**

Penz (2009) presented a detailed description of the morphology of wings and genitalia of four species of

Dasyophthalma (except *D. delanira* **stat. rest.**) and the monophyly of the “*rusina*-group” was supported by seven morphological synapomorphies: (1) dorsal forewings and hindwings with a blue iridescence below discal cell and on the submedial area, respectively; (2) dorsal forewings with yellow scales located inside cell R5 with bifid edge; (3) dorsal forewings of males without yellow markings across costal margin; (4) dorsal hindwings of males with a brown hairpencil at base of discal cell; (5) hindwings with brown androconial patch on Rs-M1; (6) ventral hindwings with postmedial band with well-defined edges; and (7) a presence of a small midline extension on the posterior portion of sterigma. Six out of the seven above synapomorphies are observed in *Dasyophthalma delanira* **stat. rest.** (the seventh was not observed since no females of *D. delanira* **stat. rest.** were available for the present study).

In addition, *D. delanira* **stat. rest.** can be distinguished from the other two species of the “*rusina*-group” by the following morphological characters: (1) dorsal forewings of males with postmedial yellow band broader than in *D. geraensis* and in *D. rusina*; and (2) dorsal hindwings with postmedial band broader than in *D. rusina*, and not as yellowish as in *D. geraensis*. The overall morphology of male genitalia is very similar to all species of the genus, but the valva has a dorsal row of spines slightly different to those illustrated by Penz (2009). However, the number and shape of the spines present high intraspecific variability, and since a single male specimen of *D. delanira* **stat. rest.** was available to study, this character was not analyzed in further details.

4.2. Conservation

The genus *Dasyophthalma* can be considered of high conservation interest for three main reasons: (1) the genus is endemic to the Atlantic Forest, with very few records in riparian forests of the Cerrado savannas, (2) three out of the five described species are of conservation interest, and (3) these are large dayflying butterflies, with a large appeal to be used in citizen science. Accordingly, getting photographic records in social media or from butterfly watchers is relatively easy, since these large butterflies call attention when present (as can be seen in this work). The geographical data is abundant and covers possibly the actual geographic distribution of all five species, allowing adequate niche modelling (except maybe for *D. delanira* **stat. rest.**), including past and future scenarios. Moreover, most species in the genus present a strong potential to be used as models in studies of molecular population ecology and phylogeography in the Atlantic Forest.

However, as previously mentioned, there are few data on natural history for all species *Dasyophthalma* (except for the two more widespread species). A good start would be describing the early stages of all species in detail (including complementing data for *D. rusina* and *D. creusa*). As for other species of Brassolini, females easily oviposit in plastic bags with leaves of the host plants; this method resulted in dozens of fertilized eggs of *D. creusa* and

D. rusina in previous attempts and larvae easily accepted leaves of the palm trees *Bactris*, *Astrocaryum* and *Euterpe* (the records on bamboos need to be confirmed) (AVLF unpublished results). Accordingly, rearing the other three species relies only on obtaining wild-caught fertilized females.

An important action would be searching for additional localities for the three threatened species in the genus (as suggested in Freitas and Marini-Filho 2011) in order to improve the calculations of AOO and EOO. For *D. delanira* **stat. rest.**, candidate areas are all in the hilltops above 1300 m in the area between Nova Friburgo, Casimiro de Abreu and Cachoeiras de Macacu (in the state of Rio de Janeiro). For *D. geraensis*, the recent records in the “Serra do Papagaio” in Alagoa municipality and Baependi municipality, both localities in the state of Minas Gerais, suggest that this species could be more widespread through the west slopes of the Mantiqueira mountain range, in some more drier and seasonal areas. Finally, for *D. vertebralis* it is very important to find any extant population within the potential geographical range; based on museum specimens, this species was last collected in 1948 (more than 80 years ago) and since then no additional individual or photo/image has been reported so far. The Sooretama Biological Reserve is the first candidate area, since the forests are virtually the same as when the two females have been collected there in 1948. Other candidate areas in the state of Espírito Santo include the lowland forests in the center of the state, in the vicinities of Santa Teresa and Santa Leopoldina and Cariacica, and some of the last larger forest remnants in the west, in region of Castelo, São Vicente and Burarama.

5. Conclusion

As much biological information about the genus *Dasyophthalma* is lacking, the present work contributes with updated data on the phylogeny, taxonomy, geographic distribution and geographic range estimates for the genus. Therefore, by including molecular data for all five species of the genus, the present study confirms the monophyly of *Dasyophthalma* and the organization of its species in two clades. In addition, data on genetic divergence supported the decision of reinstating the specific status of *D. delanira* **stat. rest.**, as the sister species of *D. geraensis* and not as a subspecies of *D. rusina*. All five species of *Dasyophthalma* are endemic to the Atlantic Forest (with *D. rusina* also present in some riparian forests in areas of Cerrado savannas adjacent to the former domain), making this an excellent model group to be studied in terms of diversification in the Atlantic Forest. In addition, two out of the five species in the genus present narrow distribution ranges and are threatened (*D. delanira* **stat. rest.** and *D. geraensis*), with huge potential to be included in monitoring programs in the protected areas where they are present. In short, by combining most of the available information, the present study can serve as a starting point for future

studies on *Dasyophthalma*, adding information that can be crucial for future conservation actions and essentially assuring the future of the threatened species in this genus.

6. Competing interests

The authors have declared that no competing interests exist.

7. Acknowledgments

We thank Mirna Casagrande, Eduardo Carneiro, Olaf Mielke for the access to material deposited in DZUP collection; Alexandre Soares for the access to material deposited in MNRJ collection; Márcio Félix, Felipe Moreira, and Claudia Rodrigues for the access to material deposited in CEIOC collection; Marcelo Duarte and Renato Silva for the access to material deposited in MZUSP collection; Thomas Pape for the access to material deposited in ZMUC; Tobias Malm for the access to material deposited in NHRS; To Carlos Guilherme Costa Mielke, Max Peters, Carlos Roberto Silva Silva and Jalmirez Silva that provided personal observation / and or photos of some live species in their habitats. To all people who provided data of *D. creusa* and *D. rusina* in iNaturalist website. To Alexandre Antonelli, Thomas Berg and RPPN Bacchus staff for sharing their place to our field work. To Junia Carrera and André Tacioli for their help in field work. To an anonymous reviewer and Dr. Ângelo Pinto for their valuable comments and suggestions to improve the manuscript; RRS was supported by FAPESP (2020/09595-0). EPB was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP (2012/03750-8, 2016/15873-8 and 2018/21432-0) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (162673/2020-5). AHBR and PAM thanks the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brazil (CAPES) – Finance Code 001. PAM thanks Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP (2021/13396-6). AVLF thanks (FAPESP) (BIOTA-FAPESP grants 2013/50297-0, 2021/03868-8) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (421248/2017-3, 304291/2020-0). This publication is part of the “RedeLep – Rede Nacional de Pesquisa e Conservação de Lepidópteros” – SISBIOTA-Brasil/CNPq (563332/2010-7). Butterfly species are registered in the SISGEN (A842AC0).

8. References

- Anisimova M, Gil M, Dufayard JF, Dessimoz C, Gascuel O (2011) Survey of branch support methods demonstrates accuracy, power, and robustness of fast likelihood-based approximation schemes. *Systematic Biology* 60(5): 685–699.
- Azevedo CO, Vaz-de-Mello FZ, Tavares MT, Brescovit AD, Marco Jr PD, Furieri KS, Brown Jr KS, Freitas AVL (2007) Capítulo 8 Os invertebrados terrestres ameaçados de extinção no estado do Espírito Santo. In: Passamani M, Mendes SL (Orgs) *Espécies da fauna ameaçadas de extinção no Estado do Espírito Santo*. Vitória: Instituto de Pesquisa da Mata Atlântica, 105–119.
- Bachman S, Moat J, Hill A, de la Torre J, Scott B (2011) Supporting Red List threat assessments with GeoCAT: geospatial conservation assessment tool. *ZooKeys* 150: 117–126. <https://doi.org/10.3897/zookeys.150.2109>
- Beccaloni GW, Vitoria AL, Hall SK, Robinson GS (2008) Catalogue of the hostplants of the Neotropical butterflies/Catálogo de las plantas huésped de las mariposas neotropicales. *Monografías Tercer Milenio*, V.8, Zaragoza, España, 536 pp.
- Brown Jr. KS (1992) Borboletas da Serra do Japi: Diversidade, hábitos, recursos alimentares e variação temporal. In: Morellato LPC (Ed.) *História natural da Serra do Japi. Ecologia e preservação de uma área florestal no Sudeste do Brasil*. Editora da Unicamp/Fapesp, Campinas, 142–187.
- Casagrande MM (2004) Brassolini. In: Heppner JB (Ed) *Atlas of Neotropical Lepidoptera. Part 4A Checklist*, Lamas G. (Ed). Association of Tropical Lepidoptera/Scientific Publishers; Gainesville, 201–205.
- Casagrande MM, Mielke OHH (2000) Larva de quinto estágio e pupa de *Dasyophthalma rusina rusina* (Godart) (Lepidoptera, Nymphalidae, Brassolinae). *Revista brasileira de Zoologia* 17: 401–404. <https://doi.org/10.1590/S0101-81752000000200011>
- Casagrande MM, Mielke OHH (2003) Larvas de quarto e quinto estágios e pupa de *Dasyophthalma creusa creusa* (Hübner) (Lepidoptera, Nymphalidae, Brassolinae). *Revista brasileira de Zoologia* 20: 157–160. <https://doi.org/10.1590/S0101-81752003000100020>
- Casagrande MM, Mielke OHH (2008) *Dasyophthalma geraensis* Rebel, 1922. In: Machado ABM, Drummond GM, Paglia AP (Orgs) *Livro Vermelho da Fauna Brasileira Ameaçada de Extinção*. Ministério do Meio Ambiente, Fundação Biodiversitas Brasília-DF, Belo Horizonte-MG, 407–408.
- Casagrande MM, Mielke OHH, Brown Jr KS (1998) Borboletas (LEPIDOPTERA) ameaçadas de extinção em Minas Gerais, Brasil. *Revista brasileira de Zoologia* 15(1): 241–259. <https://doi.org/10.1590/S0101-81751998000100021>
- COPAM (1996) Deliberação COPAM nº 041/95. Aprova a lista de espécies ameaçadas de extinção da fauna do Estado de Minas Gerais. Minas Gerais. Órgão Oficial dos Poderes do Estado. Diário Executivo, Legislativo e Publicações de Terceiros, Belo Horizonte, 104(14), parte I, 1–4.
- COPAM (2010) Deliberação normativa COPAM nº 147, de 30 de abril de 2010 – Aprova a Lista de Espécies Ameaçadas de Extinção da Fauna do Estado de Minas Gerais. Diário do Executivo do Estado de Minas Gerais, Belo Horizonte. <http://www.siam.mg.gov.br/sla/download.pdf?idNorma=13192>
- d’Araújo e Silva AG, Gonçalves CR, Galvão DM, Nascimento e Silva M, Simoni L (1968) Quarto Catálogo dos insetos que vivem nas plantas do Brasil. Ministério de Agricultura, Rio de Janeiro, 622 pp.
- deVries PJ (1987) The butterflies of Costa Rica and their natural history, vol I: Papilionidae, Pieridae, Nymphalidae. Princeton University Press, Princeton, NJ, 327 pp.
- Dudley N (2008) Guidelines for Applying Protected Area Management Categories. IUCN, Gland, Switzerland, 143 pp. <https://portals.iucn.org/library/sites/library/files/documents/pag-021.pdf>
- Espírito Santo (2005) Decreto Nº1499-R de 13 de junho de 2005 – Declara as espécies da Fauna e Flora silvestres ameaçadas de extinção no Estado do Espírito Santo e dá outras providências. Diário Oficial do Estado do Espírito Santo, Poder Executivo Vitória. <https://institutolife.org/wp-content/uploads/2018/11/Lista-da-Fauna-e-Flora-Ameacadas-de-Extincao-ES.pdf>
- Fraga CN, Peixoto AL, Leite YLR, Santos, ND, Oliveira JRPM, Sylvestre LS, Schwartzburd PB, Tuler AC, Freitas J, Lirio EJ, Couto DR, Dutra VF, Waichert C, Guerra Sobrinho T, Hostim-Silva M, Ferreira RB, Bérnills RS, Costa LP, Chaves FG, Formigoni MH, Silva JP, Ribeiro RS, Reis JC, Capellão RT, Lima RO, Saiter FZ, Lopes AS, Paglia AP, Chautems A, Braz AG, Mónico AT, Salino A, Firmino

- AD, Chagas AP, Colodetti AF, Krahel AH, Sousa AAC, Pavan ACO, Castello ACD, Loss AC, Srbeek-Araujo AC, Scudeler AL, Farro APC, Feijó A, Machado AFP, Ferreira ALN Fontana AP, Freitas AVL, Cabral A, Bolzan AAS, Weigand A, Frazão A, Neto ACR, Almeida AP, Brescovit AD, Argôlo AJS, Soares AS, Gonçalves AM, Santiago ACP, Giaretta A, Rosa AHB, Katz AM, Gomes BM, Øllgaard B, Silva BCB, Canestraro BK, Melo BF, Evaldt BHC, Amorim BS, Pimentel CR, Possamai CB, Ferreira CDM, Guidorizzi CE, Silva CBM, Ferreira CD, Zocca C, Miranda CV, Kameyama CDC, Vilar CC, Canedo C, Sampaio CA, Carneiro CE, Mynssen CM, Pessoa CS, Cunha CJ, Sá CFC, Imig DC, Sampaio D, Monteiro D, Moreira DO, Robayo DS, Machado DF, Peralta DF, Silva DF, Gonzaga DF, Zeppelini D, Silva EMP, Lozano ED, Melo E, Chiquito EA, Rossetto EFS, Smidt EC, Martins ER, Santos Neto EA, Silva Júnior EMS, Massariol FC, Firetti F, Novelli FZ, Di Dario F, Gonzatti F, Guimarães FV, Cabral FN, Mello Fraga FRM, Matos FB, Dagosta FCP, Vaz-de-Mello FZ, Souza FS, Leite FT, Molina FB, Barroso FG, Alves FM, Salles FF, Mendonça GC, Siqueira GS, Pinto GO, Romão GO, Antar GM, Rosa GAB, Zassemer G, Heiden G, Shimizu GH, Caitano HA, Lima HC, Resende HC, Pichler HA, Bergallo HG, Fernandes HQB, Secco HKC, Costa HC, Dias HM, Daros Filho HJ, Silva HL, Ornellas IS, Bergher IS, Pinheiro IEG, Koch I, Rodrigues IMC, Passamani JA, Delabie JHC, Oliveira JCF, Lubert J, Prado J, Lopes JC, Christ JA, Nodari JZ, Tonini JFR, Zorzanelli JPF, Condaack JPS, Lacerda JVA, Adelir-Alves J, Jardim JG, Santander-Neto J, Trezena JS, Schoederer JH, Gomes JML, Silva JN, Trarbach J, Rossini J, Kulkamp J, Pereira JBS, Prado JR, Guarnier JC, Souza JP, Alevis KCC, Furieri KS, Costa KG, Alves KP, Pederneiras LC, Cardoso LJT, Geise L, Mathias LB, Ingenito LFS, Merçon L, Contaifer LS, Bissoli LB, Aona LYS, Silva LTP, Freitas LT, Calazans LSB, Marinho LC, Lucas LAS, Silva EF, Lima LV, Martins LR, Kollmann LJC, Bernacci LC, Mayorga LFSP, Sarmento-Soares LM, Góes-Neto LAA, Duboc LF, Fonseca LHM F, Junior RR, Beyer M, Vianna Filho MDM, Devecchi MF, Passamani M, Britto MR, Pereira MR, Simonelli M, Trovó M, Fukuda MV, Verdi M, Pellegrini MOO, Coelho MAN, Lehnert M, Alves MAS, Kierulff MCM, Loliola MIB, Marchioreto MS, Saka MN, Rodrigues MR, Zanin M, Facco MG, Zortéa M, Freitas MO, Pastore M, Camelo MC, Milward-de-Azevedo MA, Ribeiro M, Teixeira MDR, Klautau M, Kaehler M, Menezes NA, Bigio NC, Pena NTL, Shibatta OA, Silva OLM, Gonçalves PR, Santos PM, Grossi PC, Buckup PA, Chaves PB, Paiva PC, Windisch PG, Barros PHD, Evangelista PHL, Pedro PMG, Cardoso FPH, Peloso PLV, Santos PMLA, Taucce PPG, Cardoso PCA, Almeida RF, Barbosa-Silva RG, Trad RJ, Vanstreels RET, Macieira RM, Monteiro RF, Viveros RS, Ribeiro RTM, Romanini RP, Hirai RY, Betzel RL, Asprino RC, Hurtado PR, Sousa-Lima RS, Pagotto RV, Mello-Silva R, Goldenberg R, Vicente RE, Lourenço-de-Moraes R, Bianchi RSCRC, Paresque R, Valadares RT, Guimarães RR, Ribon R, Martins-Pinheiro RF, Marquete R, Subirá RJ, Siciliano S, Recla SS, Sandra Ribeiro S, Nunes SF, Mendes SL, Oliveira TPR, Carrizo TT, SilvaTG, Volpi TA, Almeida TE, Flores TB, Kloss TG, Castro TM, Silva-Soares T, Barbosa TDM, Tavares VC, Fagundes V, Verdade VK, Amaral VS, Orrico VGD, Vale V, Junior VC, Dittrich VAO, Freitas VC, Giglio VJ, Weslei Pertel W, Colombo WD, Cardoso WC, Nóbrega YC (2019) Lista da fauna e flora ameaçadas de extinção no estado do Espírito Santo. In: Fraga CN, Formigoni MH, Chaves FG (Org) Fauna e flora ameaçadas de extinção no estado do Espírito Santo. 1ed. Instituto Nacional da Mata Atlântica, Santa Teresa, v.1, 342–397.
- Freitas AVL, Benson WW, Marini-Filho OJ, Carvalho RM (1997) Territoriality by the dawn's early light: the Neotropical owl butterfly *Caligo idomenaeus* (Nymphalidae: Brassoliniæ). *Journal of Research on the Lepidoptera* 34: 14–20. <https://doi.org/10.5962/p.266557>
- Freitas AVL, Marini-Filho OJ (2011) Plano de Ação Nacional para Conservação dos Lepidópteros Ameaçados de Extinção. Instituto Chico Mendes de Conservação da Biodiversidade, Brasília, 124pp.
- Freitas AVL, Marini-Filho OJ, Mielke OHH, Casagrande MM, Brown Jr. KS, Kaminski LA, Iserhard CA, Ribeiro, DB, Dias FMS, Dolibaina, DR, Carneiro E, Uehara-Prado M, Romanowski HP, Emery EO, Accacio GM, Rosa AHB, Bizarro JMS, Silva ARM, Guimarães MP, Silva NAP, Braga L, Almeida G. (2018a) *Dasyophthalma rusina delanira* Hewitson, 1862. In: Instituto Chico Mendes de Conservação da Biodiversidade. (Org.). Livro Vermelho da Fauna Brasileira Ameaçada de Extinção Volume VII – Invertebrados. ICMBio, Brasília, 113–115.
- Freitas AVL, Marini-Filho OJ, Mielke OHH, Casagrande MM, Brown Jr. KS, Kaminski LA, Iserhard CA, Ribeiro, DB, Dias FMS, Dolibaina, DR, Carneiro E, Uehara-Prado M, Romanowski HP, Emery EO, Accacio GM, Rosa AHB, Bizarro JMS, Silva ARM, Guimarães MP, Silva NAP, Braga L, Almeida G. (2018b) *Dasyophthalma geraensis* Rebel, 1922. In: Instituto Chico Mendes de Conservação da Biodiversidade. (Org.). Livro Vermelho da Fauna Brasileira Ameaçada de Extinção Volume VII – Invertebrados. ICMBio, Brasília, 111–113.
- Freitas AVL, Marini-Filho OJ, Mielke OHH, Casagrande MM, Brown Jr. KS, Kaminski LA, Iserhard CA, Ribeiro, DB, Dias FMS, Dolibaina, DR, Carneiro E, Uehara-Prado M, Romanowski HP, Emery EO, Accacio GM, Rosa AHB, Bizarro JMS, Silva ARM, Guimarães MP, Silva NAP, Braga L, Almeida G. (2018c) *Dasyophthalma vertebralis* Butler, 1869. In: Instituto Chico Mendes de Conservação da Biodiversidade. (Org.). Livro Vermelho da Fauna Brasileira Ameaçada de Extinção Volume VII – Invertebrados. ICMBio, Brasília, 115–117.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299.
- Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology* 59(3): 307–321. <https://doi.org/10.1093/sysbio/syq010>
- Hoffmann F (1936). Beiträge zur Naturgeschichte brasilianischer Schmetterlinge. *Entomologisches Jahrbuch* 45: 81–93.
- Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2018) UFBoot2: Improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* 35: 518–522. <https://doi.org/10.1093/molbev/msx281>
- IBAMA, 1989. Portaria IBAMA Nº 1.522, de 19 de dezembro 1989. Dispõe sobre a Lista Oficial de Espécies da Fauna Brasileira Ameaçada de Extinção. Diário Oficial da União, Brasília. DF, Seção 1: 24156.
- IUCN (2012) Guidelines for application of IUCN Red List Criteria at regional and national levels (Version 4.0). IUCN. Gland, Switzerland and Cambridge. <https://portals.iucn.org/library/sites/library/files/documents/RL-2012-002.pdf>
- IUCN Standards and Petitions Committee (2012) Guidelines for using the IUCN Red List Categories and Criteria (version 15.1). Prepared by the Standards and Petitions Committee. <https://www.iucnredlist.org/resources/redlistguidelines>

- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast Model Selection for Accurate Phylogenetic Estimates. *Nature Methods* 14: 587–589. <https://doi.org/10.1038/nmeth.4285>
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012). Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28(12): 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16: 111–120. <https://doi.org/10.1007/BF01731581>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) Mega X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution* 35: 1547–159. <https://doi.org/10.1093/molbev/msy096>
- MMA (2003) Instrução Normativa Nº 003 de 28 de maio de 2003. Diário Oficial da União, Brasília. DF, Seção 1: 88.
- MMA (2014) Lista Nacional Oficial de Espécies da Fauna Ameaçadas de Extinção-Anexo I à Portaria Nº 444, de 17 de dezembro de 2014. Diário Oficial da União, Brasília. DF, Seção 1(245): 121–126.
- MMA, 2022. Portaria MMA nº 148, de 7 de junho de 2022 Altera os Anexos da Portaria nº 443, de 17 de dezembro de 2014, da Portaria nº 444, de 17 de dezembro de 2014, e da Portaria nº 445, de 17 de dezembro de 2014, referentes à atualização da Lista Nacional de Espécies Ameaçadas de Extinção. Diário Oficial da União, Brasília. DF, Seção 1(108): 74.
- Matos-Maravi, P, Wahlberg N, Freitas AVL, deVries P, Antonelli A, Penz CM (2021) Mesoamerica is a cradle and the Atlantic Forest is a museum of Neotropical butterfly diversity: insights from the evolution and biogeography of Brassolini (Lepidoptera: Nymphalidae). *Biological Journal of the Linnean Society* 133(3): 704–724. <https://doi.org/10.1093/biolinnean/blab034>
- Meyer M, Kircher M (2010) Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harbor Protocols*, 2010(6): pdb.prot5448. <https://doi.org/10.1101/pdb.prot5448>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *Proceedings of the Gateway Computing Environments Workshop (GCE)*, New Orleans, LA, 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R (2020) IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Molecular Biology and Evolution* 37(5): 1530–1534. <https://doi.org/10.1093/molbev/msaa015>
- Nogueira TA (2012) Análise da diversidade e efeito de borda na assembléia de borboletas frugívoras da Reserva Biológica de Sooretama–ES. Master degree Thesis Universidade Federal do Espírito Santo, Vitória, Espírito Santo, Brazil.
- Otero LS, Brown Jr. KS, Mielke OHH, Monteiro RF, Costa JM, Macêdo MV, Maciel NC, Becker J, Salgado NC, Santos SB, Moya GE, Almeida JM, Silva MD (2000) Invertebrados terrestres. In: Bergallo GH, Rocha CFD, Sluys MV, Alves MAS (Orgs) *A fauna ameaçada de extinção do estado do Rio de Janeiro*. Editora UERJ, Rio de Janeiro, 53–62.
- Penz CM (2007) Evaluating the monophyly and phylogenetic relationships of Brassolini genera (Lepidoptera, Nymphalidae). *Systematic Entomology* 32: 668–689. <https://doi.org/10.1111/j.1365-3113.2007.00391.x>
- Penz CM (2009). Phylogeny of *Dasyophthalma* butterflies (Lepidoptera, Nymphalidae, Brassolini). *Insecta Mundi* 69: 1–12. <http://doi.org/10.5281/zenodo.5405005>
- Rosa AHB, Ribeiro DB, Freitas AVL (2023) How data curation and new geographical records can change the conservation status of threatened Brazilian butterflies. *Journal of Insect Conservation*, online version. <https://doi.org/10.1007/s10841-023-00464-0>
- Santos JP, Freitas AVL, Brown KS Jr., Carreira JYO, Gueratto PE, Rosa AHB, Lourenço GM, Accacio GM, Uehara-Prado M, Iserhard CA, Richter A, Gawlinski K, Romanowski HP, Mega NO, Teixeira MO, Moser A, Ribeiro DB, Araujo PF, Filgueiras BKC, Melo, DHA, Leal IR, Beirão MV, Ribeiro SP, Cambuí ECB, Vasconcelos RN, Cardoso MZ, Paluch M, Greve RR, Voltolini JC, Galetti M, Regolin AL, Sobral-Souza T, Ribeiro MC (2018) Atlantic butterflies: a data set of fruit-feeding butterfly communities from the Atlantic forests. *Ecology* 99(12): 2875–2875. <https://doi.org/10.1002/ecy.2507>
- São Paulo (2008) Decreto nº 53.494, de 02 de outubro de 2008. Declara as Espécies da Fauna Silvestre Ameaçadas, as Quase Ameaçadas, as Colapsadas, Sobreexplotadas, Ameaçadas de Sobreexplotação e com dados insuficientes para avaliação no Estado de São Paulo e dá providências correlatas. Diário Oficial Poder Executivo – Seção I, 118(187): 1.
- São Paulo (2010) Decreto nº 56.031, de 20 de julho de 2010 – Declara as Espécies da Fauna Silvestre Ameaçadas, as Quase Ameaçadas, as Colapsadas, Sobreexplotadas, Ameaçadas de Sobreexplotação e com dados insuficientes para avaliação no Estado de São Paulo e dá providências correlatas. Diário Oficial Poder Executivo – Seção I, São Paulo, 120(136): 3.
- São Paulo (2014) Decreto nº 60.133, de 7 de fevereiro de 2014 – Declara as espécies da fauna silvestre ameaçadas de extinção, as quase ameaçadas e as deficientes de dados para avaliação no Estado de São Paulo e dá providências correlatas. Diário Oficial Poder Executivo – Seção I, São Paulo, 124(27): 25.
- São Paulo (2018) Decreto nº 63.853, de 27 de novembro de 2018 – Declara as espécies da fauna silvestre no Estado de São Paulo regionalmente extintas, as ameaçadas de extinção, as quase ameaçadas e as com dados insuficientes para avaliação, e dá providências correlatas. Diário Oficial Poder Executivo – Seção I, São Paulo, 128 (220–221): 1.
- Schmith K, Hoffmann F (1931) Die Brassoliden des Municip Joinville im Staate Sta. Catharina, Südbrasilien. *Entomologisches Jahrbuch* 40: 150–153.
- SEMA (1998) Portaria SEMA nº 01, de 04 de junho de 1998 – Dispõe sobre a pesquisa realizada pela Universidade do Estado do Rio de Janeiro – UERJ, elaborando a Lista da Fauna Ameaçada de Extinção no Estado. Diário Oficial do Estado do Rio de Janeiro, Poder Executivo, parte 1, 102: 9–16.
- Shirai LT, Barbosa EP, Greve RR, Magaldi LM, Nascimento AR, Freitas, AVL (2017) Natural history of *Selenophanes cassiope guarany* (Lepidoptera: Nymphalidae: Brassolini): an integrative approach, from molecules to ecology. *Annals of the Entomological Society of America*, 110(2): 145–159. <https://doi.org/10.1093/aesa/saw068>
- Shirai LT, Machado PA, Mota LL, Rosa AHB, Freitas AVL (2019) DnB., the Database of Nymphalids in Brazil., with a Checklist for Standardized Species Lists. *The Journal of the Lepidopterists' Society* 73(2): 93–108. <https://doi.org/10.18473/lepi.73i2.a4>
- Silva-Brandão KL, Freitas AVL, Brower AVZ, Solferini VN (2005) Phylogenetic relationships of the New World Troidini swallowtails

- (Lepidoptera: Papilionidae) based on COI, COII, and EF-1 α genes. Molecular Phylogenetics and Evolution 36(3): 468–483. <https://doi.org/10.1016/j.ympev.2005.04.007>
- Twort VG, Minet J, Wheat CW, Wahlberg N (2021) Museomics of a rare taxon: placing Whalleyanidae in the Lepidoptera Tree of Life. Systematic Entomology 46: 926–937. <https://doi.org/10.1111/syen.12503>
- Zikán JF, Zikán W (1940). Introdução para o catálogo da insetofauna do Itatiaia e da Mantiqueira. Rodriguésia (Rio de Janeiro) 4(13): 155–165.

Supplementary Material 1

Figure S1, Table S1

Authors: Rosa AHB, Barbosa EP, Machado PA, Siewert RR, Freitas AVL (2023)

Data type: .docx

Explanation note: **Figure S1.** Adults and habitat of *Dasyophthalma vertebralis* and *D. creusa* from Sooretama Biological Reserve (SBR), Sooretama, Espírito Santo state. **A, B** The two *D. vertebralis* female specimens from CEIOC collection; **C, D** two *D. creusa* female specimens from CEIOC collection; **E** closer view of *D. vertebralis* and *D. creusa* habitat on a road of SBR; **F** attractive traps in the interior of forest of SBR. — **Table S1.** Abundance data of *D. creusa* and *D. rusina*.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/asp.81.e96397.suppl1>

Supplementary Material 2

Concatenated dataset with aligned sequences

Authors: Rosa AHB, Barbosa EP, Machado PA, Siewert RR, Freitas AVL (2023)

Data type: .phy

Explanation note: Concatenated dataset with aligned sequences.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/asp.81.e96397.suppl2>